

Available online at [www.sciencerepository.org](http://www.sciencerepository.org)

Science Repository



## Short Communication

# Duration and rate of clinical eruption of third molars

Ashraf I. Shaweesh<sup>1,2\*</sup>

<sup>1</sup>Senior lecturer in Oral Health Therapy, School of Dentistry and Health Sciences, Faculty of Science, Charles Sturt University, Wagga Wagga, New South Wales, 2678, Australia

<sup>2</sup>Professor of Oral Science, Faculty of Dentistry, Jordan University of Science and Technology, PO Box 3030, Ramtha, Irbid, 22110, Jordan

### ARTICLE INFO

#### Article history:

Received: 14 January, 2019

Accepted: 8 February, 2019

Published: 26 February, 2019

#### Keywords:

Clinical eruption

duration of eruption

eruption rate

third molars

### ABSTRACT

**Objectives:** The aim of the current paper was to present norms of clinical duration and rate of eruption for upper and lower 3<sup>rd</sup> molars utilizing recently published data for a Jordanian population.

**Methods:** Duration of Clinical Eruption (CDE) in years and Clinical Eruption Rates (CER) in mm/year were calculated from recently published median ages at emergence and functional eruption of 3<sup>rd</sup> molars for a Jordanian population.

**Results:** Third molars clinically erupt at a rate between 1.45 to 1.85 mm/year, being slightly faster in the lower jaw. The average CDEs were 2.95 and 2.75 years for upper and lower 3<sup>rd</sup> molars respectively. Upper 3<sup>rd</sup> molars of males needed a slightly longer CDE than lower 3<sup>rd</sup> molars did while, in females, there was no difference in CDE across arches. The average CERs were 1.53 and 1.82 mm/year for upper and lower 3<sup>rd</sup> molars respectively. In contrast to males, females exhibited slightly faster CER for the upper 3<sup>rd</sup> molar and slightly slower CER for the lower 3<sup>rd</sup> molar. CERs are steadier and more linear in the upper than in the lower 3<sup>rd</sup> molar.

**Conclusions:** This study has reported the first norms of duration and rate of clinical eruption specific to upper and lower 3<sup>rd</sup> molars. Normally, third molars remain around 3 years in clinical eruption, which render them more vulnerable to recurrent pericoronitis and other local pathologies than the rest of the teeth. The availability of these norms will help dentists and auxiliaries identify and diagnose delayed and slow eruption cases and make their clinical decision upon removing or keeping 3<sup>rd</sup> molars.

© 2019 Ashraf I. Hosting by Science Repository. All rights reserved.

## Introduction

Although timing of 3<sup>rd</sup> molar eruption is genetically determined, it can be affected by a variety of developmental, environmental and iatrogenic factors such as insufficient jaw space, malalignment of other teeth, premolar / first molar extraction for orthodontic treatment, second molar extraction, rapid maxillary expansion and loss of mesio-distal arch space due to caries [1-10]. Since 3<sup>rd</sup> molars are the last permanent teeth to emerge in the arch, they are more commonly vulnerable to delayed eruption and malalignment than other teeth, particularly when jaw space is insufficient. This leads to a higher chance for the development of

localized pathologies such as impaction, crowding and malalignment of other teeth, cystic changes, pathologic tooth resorption, pericoronitis and periodontitis [11-13].

The duration of time a 3<sup>rd</sup> molar needs from gingival emergence to functional eruption and the clinical rate of its eruption are very significant, and ideally, should be fast enough to reduce the frequency of recurrent episodes of pericoronitis and reduce the chance of occurrence of other pathologies associated with long-standing partially-erupting 3<sup>rd</sup> molars. Moreover, dentists should be able to periodically assess whether an emerged 3<sup>rd</sup> molar is clinically erupting at a normal eruption rate

\*Correspondence to: Ashraf I. Shaweesh, Senior lecturer in Oral Health Therapy, School of Dentistry and Health Sciences, Faculty of Science, Charles Sturt University, Wagga Wagga, New South Wales, 2678, Australia; Tel: +61 2 6933 2673; E-mail: [ashaweesh@csu.edu.au](mailto:ashaweesh@csu.edu.au)

before making any decision upon removing or keeping the tooth. In this regard, population-specific norms on clinical duration and rate of 3<sup>rd</sup> molar eruption would be very useful guides.

A very limited number of studies have been published about clinical eruption rates of permanent teeth and most of them are dated [14-17]. To the best of the author's knowledge, and apart from only a 40-year-old paper presenting the eruption rate of upper 3<sup>rd</sup> molars, the dental literature is devoid of any publications specifically reporting the eruption rate of upper and lower 3<sup>rd</sup> molars [18]. The aim of the current paper was to present norms of clinical duration and rate of eruption of upper and lower 3<sup>rd</sup> molars utilizing recently published data on a Jordanian population.

## Materials and Methods

The current work is part of an extended project titled "chronology of eruption of permanent dentition in the Jordanians" that received the ethics approval from the Deanship of Research at Jordan University of Science and Technology (ID# 13/2007). Several data on timing of permanent tooth eruption and its correlates have been published out of this project including the one on timing of third molar eruption [19]. The current study used the data on eruption timing of third molars to investigate the rate and clinical duration of eruption of third molars in a Jordanian population.

The published data on the timing of eruption of upper and lower 3<sup>rd</sup> molars were derived from a sample of 571 Jordanian adolescents and young adults aged 15 – 27 years (275 males and 296 females) [19]. All participants provided signed consent to take part in the study. None of the participants had presented any clinical signs of 3<sup>rd</sup> molar impaction or agenesis and those who had showed radiographic evidence were excluded. In addition, participants confirmed to have had skeletal / growth disorders, obesity, moderate to severe crowding, spacing and tooth malalignment or received oral or jaw surgeries, trauma or orthodontic treatment had been excluded as well [19]. Probit regression analysis was used to calculate the median ages at which upper and lower 3<sup>rd</sup> molars reach four clinical eruption stages (Gingival Emergence - GEM, Occlusal Surface Eruption - OSE, Half Crown Eruption - HCE and Full Functional Eruption - FFE).

In the current study, Clinical Duration of Eruption (CDE), which is the duration of time while a tooth is being clinically visible commencing from gingival emergence up until full functional eruption, was calculated by subtracting the median age at GEM from the median age at FFE [20]. The Clinical Eruption Rate (CER) was given when the height of the clinical crown (the part of the tooth that is clinically visible) was divided by CDE. The average heights of the anatomical crown (the part of the tooth covered with enamel) in the upper and lower 3<sup>rd</sup> molars are 6.5 and 7.0 mm respectively [21]. Considering the average healthy gingival sulcus depth of 2.0 mm, this gives clinical crown heights of 4.5 and 5.0 mm for the upper and lower 3<sup>rd</sup> molars respectively [22]. Therefore, CERs for the upper and lower 3<sup>rd</sup> molars were calculated accordingly and expressed in the unit of mm/year.

The full clinical eruption from GEM to FFE could also be divided further into three segments; the first segment marks the eruption of the occlusal ¼ of the clinical crown height (GEM to OSE = 1.125 mm or 1.25 mm

for upper and lower 3<sup>rd</sup> molars respectively), the second marks the eruption of the second ¼ (OSE to HCE = 1.125 mm or 1.25 mm for upper and lower 3<sup>rd</sup> molars respectively) and the 3<sup>rd</sup> marks the eruption of the cervical ½ of the clinical crown height (HCE to FFE = 2.25 mm or 2.5 mm for upper and lower 3<sup>rd</sup> molars respectively). Consequently, CDEs and CERs were calculated per each of the clinical crown segment accordingly.

## Results

(Table 1) presents the CDEs in years and the CERs in mm/year for the upper and lower 3<sup>rd</sup> molars, and for males and females. (Table 1) also present the inter-arch and gender differences. In males, full CDEs were 3.10 and 2.70 years for upper and lower 3<sup>rd</sup> molars respectively, with an inter-arch difference of 0.40 years. In females, CDE for lower 3<sup>rd</sup> molar was close to that in males (2.80 years) although in the upper 3<sup>rd</sup> molar it was slightly shorter (2.80 years). In other words, upper 3<sup>rd</sup> molars of males needed a slightly longer duration of time for clinical eruption than lower 3<sup>rd</sup> molars did while, in females, there was no difference in clinical duration of eruption across arches. Regardless of gender, the average CDEs were 2.95 and 2.75 years for upper and lower 3<sup>rd</sup> molars respectively.

The rates of clinical eruption were slightly slower in the upper 3<sup>rd</sup> molar (1.45 and 1.61 mm/yrs. in males and females respectively) than in the lower (1.85 and 1.79 mm/yrs. for males and females respectively). In contrast to males, females exhibited slightly faster CER for the upper 3<sup>rd</sup> molar (1.61 vs. 1.45 mm/yrs.) and slightly slower CER for the lower 3<sup>rd</sup> molar (1.79 vs. 1.85 mm/yrs.). Regardless of gender, the average CERs were 1.53 and 1.82 mm/year for upper and lower 3<sup>rd</sup> molars respectively.

Since none of the gender and inter-arch differences in the median ages of GEM and FFE of upper and lower 3<sup>rd</sup> molars were statistically significance, it is assumed that such small differences in CDE and CER across arches and genders are non-statistically significant as well [19].

(Table 2) presents the segmented CDEs and CERs for both sexes and arches. It is generally noticed that CERs are steadier and more linear in the upper than in the lower 3<sup>rd</sup> molar with the CER of the 2<sup>nd</sup> occlusal ¼ of upper 3<sup>rd</sup> molar being the fastest somehow. In contrast lower 3<sup>rd</sup> molars showed faster CER during the eruption of the cervical ¾ of the clinical crown and slightly slower CER during the eruption of the occlusal ¼.

## Discussion

To the best of the author's knowledge, this paper has presented the first standards for duration and rate of clinical eruption of upper and lower 3<sup>rd</sup> molars. The term "clinical" was used in the context to designate the eruption period during which the tooth is "clinically" visible in the mouth [19]. Since the term "eruption" alone would also include the intra-osseous stages of tooth movements, it was more appropriate to use the term "clinical eruption" to avoid any confusions.

The current study followed a different approach to what other studies about tooth clinical eruption rates have followed. While Burke & Newell devised a photographic method for measuring active clinical eruption rates of maxillary central incisors, Berkovitz & Bass and Smith observed dental casts sequentially over a period of time and measured clinical

eruption of permanent teeth manually [16-18]. The longitudinal approach and the need to take photographs or take impressions and make dental casts limited the number of subjects studied. In the present study, the duration and rate of tooth eruption were provided upon utilizing recently published standards on median ages at different clinical eruption stages, which were calculated by clinical examination of a large number of participants.

The precision in determining the clinical eruption stage at which a given tooth presents at examination constitutes an important point to address here [19]. Obviously, there were no problems in recording GEM and FEE stages as these are readily identified upon clinical examination. GEM marked the first moment of gingival penetration and FEE the occlusal contact or functional eruption, which could be confirmed by biting on an articulating paper when the opposing tooth was present or when the tooth's occlusal surface was level with the occlusal plane taking to consideration the curves of Spree and Wilson. There might have been a greater challenge in detecting OSE and HCE at the same level of precision compared to GEM and FEE since these were determined subjectively by the examiner according to the criteria described by Shaweesh [19]. Nevertheless, the calculated clinical eruption rates in this paper relied on the median ages at reaching GEM and FEE (Table 1), which assumes linearity and steadiness in eruption rate. The median ages at reaching OSE and HCE were only used to describe the slight difference in linearity of clinical eruption rates between upper and lower 3<sup>rd</sup> molars (Table 2).

Clinical eruption rates of 3<sup>rd</sup> molars were calculated in reference to the average clinical crown height reported in Dental Anatomy textbooks [21]. In the absence of norms on occluso-cervical crown diameters of 3<sup>rd</sup> molars of Jordanians, it was necessary to use the ones reported by Nelson & Ash Jr and assume that Jordanians were not significantly different in this regard from the populations Nelson & Ash Jr reported the norms on.

Shaweesh paid all efforts to ascertain the normality of clinical eruption among the subjects examined [19]. All suspicious cases of delayed eruption due to crowding, malalignment of 3<sup>rd</sup> molars and other adjacent teeth, malocclusion and other relevant pathologies and cases of

accelerated eruption due to excessive spacing or obesity were all excluded. Besides, Shaweesh excluded potential subjects who reported receiving orthodontic treatment, which is anticipated to affect the eruption timing and clinical eruption rates of 3<sup>rd</sup> molars [1-4, 23]. Moreover, in order not to mistaken cases of 3<sup>rd</sup> molar impaction or agenesis for yet-to-erupt 3<sup>rd</sup> molars, such cases were excluded as well.

In the present paper, the CDEs of upper and lower 3<sup>rd</sup> molars were 2.95 and 2.75 years on average, which are significantly longer than those for other permanent teeth of the same population [20]. According to Shaweesh, the permanent tooth with the longest CDE was the lower second molar which needed one year on average during clinical eruption. This was followed by upper first premolar, upper canine, lower canine and upper second molar with CDEs ranging approximately from 0.94 down to 0.72 years, respectively. It is obvious in the present paper that the clinical duration of eruption in a 3<sup>rd</sup> molar is at least three folds those of other permanent teeth with long CDEs. Interestingly, Shaweesh reported an average CDE of nearly 0.18 years for first permanent molars rendering the CDE of 3<sup>rd</sup> molar nearly 15 times longer [20].

Population-specific data on CDEs for 3<sup>rd</sup> molars have not been published purposefully. However, in few studies mean / median age at FEEs and GEMs have been provided for some populations [24-28]. Consequently, CDEs could be generated from these studies and compared to the CDEs of Jordanians (Table 3). It is apparent in Table 3 that CDEs of Jordanians are generally longer than those in other populations although CDE of upper 3<sup>rd</sup> molar in German females was longer than its corresponding Jordanian CDE and the CDEs of 3<sup>rd</sup> molars in the First Nation People of Canada (except in upper 3<sup>rd</sup> molars of males) were comparable to the corresponding ones in the Jordanians [27, 28]. The Japanese and to a lesser extent Northern Chinese showed the shortest CDEs [25, 26]. Generally, except for Jordanians (current study), Croatians and Japanese, CDEs were not consistent between sexes and across arches [24]. In fact, it is not guaranteed whether these studies have followed the same strict exclusion criteria followed by Shaweesh which may explain their different CDEs compared to Jordanians and the inconsistent CDEs between sexes and across arches.

**Table 1:** Clinical Duration of Eruption (CDE) in years and Clinical Eruption Rate (CER) in mm/year for the upper and lower 3<sup>rd</sup> molars, and for males and females. Inter-gender differences in CDE and CER are also presented.

	Upper 3 <sup>rd</sup> Molar		Lower 3 <sup>rd</sup> Molar		Inter-arch difference	
	CDE	CER	CDE	CER	CDE	CER
Males	3.10	1.45	2.70	1.85	0.40	-0.40
Females	2.80	1.61	2.80	1.79	0.00	-0.18
Gender difference	0.30	-0.16	-0.10	0.06		

**Table 2:** Clinical Duration of Eruption (CDE) in years and Clinical Eruption Rate (CER) in mm/year per eruption segment, for upper and lower 3<sup>rd</sup> molars, and for males and females.

Eruption segment	Distance erupted (mm)	Upper 3 <sup>rd</sup> Molar			Lower 3 <sup>rd</sup> Molar		
		GEM - OSE	OSE - HCE	HCE - FEE	GEM - OSE	OSE - HCE	HCE - FEE
		1.125	1.125	2.25	1.25	1.25	2.50
Males	CDE	0.90	0.70	1.50	1.10	0.50	1.10
	CER	1.25	1.61	1.50	1.14	2.50	2.27
Females	CDE	0.80	0.60	1.40	1.20	0.60	1.00
	CER	1.41	1.88	1.61	1.04	2.08	2.50

GEM: Gingival Emergence, OSE: Occlusal Surface Eruption, HCE: Half Crown Eruption. FEE: Full Functional Eruption.

**Table 3:** CDEs (in years) in Jordanians and other populations.

Population	Males	Females	Males	Females
	Upper 3rd molar	Lower 3rd molar	Upper 3rd molar	Lower 3rd molar
Jordanians	3.1	2.7	2.8	2.8
Northern Chinese	2.1	0.8	1.1	1.4
Japanese	0.9	1.2	1.1	1.0
Germans	1.3	1.0	3.7	1.7
First Nation People of Canada	0.6	3.0	2.3	2.8
Croatians	1.7	1.4	1.7	1.6

On average, upper and lower 3<sup>rd</sup> molars showed CERs of 1.53 and 1.82 mm/year respectively, with the rate of lower 3<sup>rd</sup> molars being relatively faster. However, this was not consistent between sexes. Moreover, upper 3<sup>rd</sup> molars showed steadier rate of clinical eruption across segments compared to lower 3<sup>rd</sup> molars. The latter showed slower eruption of occlusal ¼ and faster eruption of the cervical ¾ of the clinical crown. In contrast to upper, lower 3<sup>rd</sup> molars may possibly need to cut through much denser overlying bone and mucosa first to clear off the occlusal surface completely from the overlying soft-tissue before they would proceed occlusally more passively. However, this topic prompts for further investigations.

Berkovitz & Bass reported a maximum clinical eruption rate in upper 3<sup>rd</sup> molars of 1 mm/2 months, which is much faster than the corresponding rate reported in the present study [18]. However, Berkovitz & Bass also reported a much slower rate of less than 1 mm/6 months which is consistent with the corresponding rate shown here. The different approach, populations and sample size and the fact that Berkovitz & Bass did not seem to have followed any exclusion criteria must be taken to consideration in comparing the results. In the absence of literature on clinical eruption rates of lower 3<sup>rd</sup> molars, only those of upper 3<sup>rd</sup> molars have been discussed in light of relevant literature.

## Conclusions

This study has purposefully reported the first norms of duration and rate of clinical eruption specific to upper and lower 3<sup>rd</sup> molars. Third molars clinically erupt at a rate between 1.45 to 1.85 mm/year, being slightly faster in the lower jaw. Normally, third molars remain around 3 years in clinical eruption, which render them more vulnerable to recurrent pericoronitis and other local pathologies than the rest of the teeth. The availability of these norms will help dentists and dental auxiliaries identify and diagnose delayed and slow eruption cases and make their clinical decision upon removing or keeping 3<sup>rd</sup> molar.

## Acknowledgment

The authors feel grateful to Charles Sturt University - Australia for providing all academic facilities in data analysis, statistical analysis and the writing up of this paper.

## Funding

The authors would like to acknowledge and thank the Deanship of Research at Jordan University of Science and Technology for funding this research project (grant number 113/2007).

## REFERENCES

- Moffitt AH (1998) Eruption and function of maxillary third molars after extraction of second molars. *Angle Orthodontist* 68: 147-152. [\[Crossref\]](#)
- Huang GJ (2007) Early mandibular first molar removal because of caries facilitates eruption of the ipsilateral third molar. *J Evid Based Dent Prac* 7: 25-26. [\[Crossref\]](#)
- Salehi P, Danaie SM (2008) Lower third molar eruption following orthodontic treatment. *East Mediterr Health J* 14: 1452-1458. [\[Crossref\]](#)
- Bayram M, Oezer M, Arici S (2009) Effects of first molar extraction on third molar angulation and eruption space. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 107: 14-20. [\[Crossref\]](#)
- Orton-Gibbs S, Orton S, Orton H (2001) Eruption of third permanent molars after the extraction of second permanent molars. Part 2: Functional occlusion and periodontal status. *Am J Orthod Dentofacial Orthop* 11: 239-244. [\[Crossref\]](#)
- Orton-Gibbs S, Crow V, Orton HS (2001) Eruption of third permanent molars after the extraction of second permanent molars. Part 1: Assessment of third molar position and size. *Am J Orthod Dentofacial Orthop* 119: 226-238. [\[Crossref\]](#)
- Gooris CGM, Artun J, Joondeph DR (1990) Eruption of mandibular 3rd molars after 2nd-molar extractions - a radiographic study. *Am J Orthod Dentofacial Orthop* 98: 161-167. [\[Crossref\]](#)
- Sokucu O, Ozturk F, Babacan H, Bicakci AA (2008) Does rapid maxillary expansion affect the eruption of upper third molars? *Angle Orthod* 78: 195-200. [\[Crossref\]](#)
- Chang SW, Shin SY, Kum KY, Hong J (2009) Correlation study between distal caries in the mandibular second molar and the eruption status of the mandibular third molar in the Korean population. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 108: 838-843. [\[Crossref\]](#)
- Lee M-H, Seol J-E, Jang I-G, Hong J, Kim C-S (2008) Correlation of distal caries in the mandibular second molar and eruption state of the mandibular third molar. *J Korean Associ Oral Maxillofacial Surge* 34: 640-643.
- Garcia RI, Chauncey HH (1989) The eruption of third molars in adults: a 10-year longitudinal study. *Oral Surg Oral Med Oral Pathol* 68: 9-13. [\[Crossref\]](#)
- Garcia RI, Sinclair C, Chauncey HH (1986) Internal resorption of an unerupted third molar. *Spec Care Dentist* 6: 205-207. [\[Crossref\]](#)
- Laskin DM (1971) Evaluation of third molar problem. *J Am Dent Assoc* 82: 824-828. [\[Crossref\]](#)
- Jedlinska W (1976) Variation of eruption rates of permanent teeth a longitudinal study of dental development of wroclaw poland school children. *Studies in Physical Anthropology* 1976: 85-95.
- Lemos Rinaldi MR, de Lima EM, de Menezes LM, Deon Rizzato SM, Baccarin Matje PR, et al. (2017) Eruption rates of lower second premolars at different development stages evaluated with cone-beam computed tomography. *Angle Orthod* 87: 570-575. [\[Crossref\]](#)

16. Smith RG (1980) A clinical-study into the rate of eruption of some human permanent teeth. *Arch Oral Biol* 25: 675-681. [[Crossref](#)]
17. Burke PH, Newell DJ (1958) A photographic method of measuring eruption of certain human teeth. *Am J of Orthod* 44: 590-602.
18. Berkovitz BKB, Bass TP (1976) Eruption rates of human upper 3rd molars. *J Dent Res* 55: 460-464. [[Crossref](#)]
19. Shaweesh AI (2016) Timing of clinical eruption of third molars in a Jordanian population. *Arch Oral Biol* 72: 157-163. [[Crossref](#)]
20. Shaweesh AI (2012) Clinical duration of permanent tooth eruption in Jordanians From emergence to functional eruption. *Int J Stomatology Occlusion Med* 5: 70-76.
21. Nelson S, Ash Jr. MM (2010) Wheeler's Dental Anatomy, Physiology and Occlusion. 9th ed: Saunders Elsevier.
22. Newman MG, Takei HH, Klokkevold PR, Carranza FA (2015) Carranza's clinical periodontology. 12th ed: St. Louis, MO : Saunders Elsevier.
23. Brezulier D, Fau V, Sorel O (2017) Influence of orthodontic premolar extraction therapy on the eruption of the third molars: A systematic review of the literature. *J Am Dent Assoc* 148: 903-912. [[Crossref](#)]
24. Brkic H, Vodanovic M, Dumancic J, Lovric Z, Cukovic-Bagic I, et al. (2011) The Chronology of Third Molar Eruption in the Croatian Population. *Coll Antropol* 35: 353-357. [[Crossref](#)]
25. Guo Y-c, Yan C-x, Lin X-w, Zhou H, Pan F, et al. (2014) Studies of the chronological course of third molars eruption in a northern Chinese population. *Arch Oral Biol* 59: 906-911. [[Crossref](#)]
26. Olze A, Ishikawa T, Zhu BL, Schulz R, Heinecke A, et al. (2007) Studies of the chronological course of wisdom tooth eruption in a Japanese population. *J Forensic Sci Int* 174: 203-206. [[Crossref](#)]
27. Olze A, Peschke C, Schulz R, Schmeling A (2008) Studies of the chronological course of wisdom tooth eruption in a German population. *J Forensic Leg Med* 15: 426-429. [[Crossref](#)]
28. Schmeling A, Olze A, Pynn BR, Kraul V, Schulz R, et al. (2010) Dental age estimation based on third molar eruption in First Nation people of Canada. *J Forensic Odontostomatol* 28: 32-38. [[Crossref](#)]