

Identification of Beak and Feather Disease Virus in an Unusual Novel Host (*Merops ornatus*) Using Nested PCR

Subir Sarker,^a Shubhagata Das,^b Karla Helbig,^a Shane R. Raidal^b

Department of Physiology, Anatomy and Microbiology, School of Life Sciences, La Trobe University, Melbourne, Victoria, Australia^a; School of Animal and Veterinary Sciences, Faculty of Science, Charles Sturt University, Wagga Wagga, New South Wales, Australia^b

The complete genome sequence of beak and feather disease virus (BFDV) was discovered from a rainbow bee-eater (*Merops ornatus*), a species of *Coraciiformes*. The genome consisted of 1,996 bp encoding two major bidirectional transcribed open reading frames. This is the first evidence of BFDV infection and complete genome characterization for this novel host species.

Received 13 October 2016 Accepted 14 October 2016 Published 8 December 2016

Citation Sarker S, Das S, Helbig K, Raidal SR. 2016. Identification of beak and feather disease virus in an unusual novel host (*Merops ornatus*) using nested PCR. *Genome Announc* 4(6):e01376-16. doi:10.1128/genomeA.01376-16.

Copyright © 2016 Sarker et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Subir Sarker, s.sarker@latrobe.edu.au.

Circoviruses are among the smallest and simplest of all known autonomously replicating viruses with a nonenveloped, icosahedral, and single-stranded circular DNA (ssDNA) genome of approximately 2 kb (1). Circovirus sequences have been identified in numerous bird species, including parrots, pigeons, gulls, ducks, geese, swans, ravens, canaries, finches, starlings, and chickens (2). To date, a wide range of psittacine species both in wild and captive populations have been documented to be infected with beak and feather disease virus (BFDV) (3–9). However, the disease status of one of the Australian migratory birds, the rainbow bee-eater, which is regularly moving to and from Asia through the Terries Strait Islands, is unknown. It has been noted that members of the family *Meropidae* (commonly bee-eaters) have had an extremely rare chance of receiving avian influenza (10). Here, we report the identification and molecular characterization of a BFDV genome in rainbow bee-eater (*Merops ornatus*), a species of *Coraciiformes* unrelated to parrots and not previously known to be susceptible to any avian circovirus.

Dried blood spots collected from a juvenile rainbow bee-eater (sample ID, 14-1186/01; year of sampling, 2014; GPS location, –23.420757°S 133.5250201°E) were used as a source of genomic DNA, and extraction of DNA was carried out using established protocols (11). Published circovirus genome sequences were retrieved from the literature and NCBI GenBank (12), and two sets of degenerate primers were designed. The nested PCR was standardized and used for the amplification of a segment of a circovirus sequence described by Sarker et al. (13). Direct Sanger sequencing of the purified gel bands resulted in a 350-bp sequence after trimming off primer sequences, and a BLASTn search with these sequences returned multiple hits to the BFDV replication-associated gene. Consequently, additional standardized PCR protocol was used for the amplification and sequencing of the remainder of the genome of BFDV (4, 6, 13).

A preliminary BLASTn (14) analysis of the assembled sequence showed a significant match to BFDV. Further analysis of the complete genome of BFDV (1,996 nucleotides) from *Merops ornatus* revealed an overall 82.2 to 95.6% pairwise identity with other

BFDV genomes available in GenBank, and the highest similarity (95.6%) to a BFDV from a wild red-tailed black cockatoo from Western Australia (KF385399) (6). The genome has the same basic structure as other BFDV genomes, which includes two major open reading frames (ORFs): ORF1 on the virion strand, encoding a replication-associated protein (Rep), and ORF2, encoding the capsid protein (Cap). Consequently, based on BLASTn (14) and BLASTp (15) analyses, ORF2 was more diverse than ORF1.

This study highlights the evidence of the rainbow bee-eater being an unusual host for BFDV infection. Some aspects of this case are difficult to explain fully without conducting ethically debatable experimental virus-transmission experiments. The rainbow bee-eater is a migratory bird that does not intimately share any ecological niche with any known parrot species. Therefore, this study may provide a unique opportunity for the better understanding of BFDV infectivity in the ecosystem.

Accession number(s). This whole-genome sequence of BFDV has been deposited in DDBJ/ENA/GenBank under the accession number [KM823541](https://www.ncbi.nlm.nih.gov/nuccore/KM823541). The version described in this paper is the first version.

ACKNOWLEDGMENTS

We acknowledge the Australian Research Council's Discovery Projects funding scheme for their financial support (grant number DP1095408). We are extremely grateful to the La Trobe University School of Life Sciences Publication Booster Award for financial support to S.S.

FUNDING INFORMATION

We acknowledge the Australian Research Council's Discovery Projects funding scheme for their financial support (grant number DP1095408). We are extremely grateful for financial support from the La Trobe University School of Life Sciences Publication Booster Award to S.S.

REFERENCES

- Sarker S, Terrón MC, Khandokar Y, Aragão A, Hardy JM, Radjainia M, Zaragoza MJ, de Pablo PJ, Coulbaly F, Buzo DL, Raidal SR, Forwood JK. 2016. Structural insights into the assembly and regulation of distinct

- viral capsid complexes. *Nat J Commun* 7:13014. <http://dx.doi.org/10.1038/ncomms13014>.
2. Delwart E, Li L. 2012. Rapidly expanding genetic diversity and host range of the *Circoviridae* viral family and other rep encoding small circular ss-DNA genomes. *Virus Res* 164:114–121. <http://dx.doi.org/10.1016/j.virusres.2011.11.021>.
 3. Raidal SR, McElnea CL, Cross GM. 1993. Seroprevalence of psittacine beak and feather disease in wild psittacine birds in New South Wales. *Aust Vet J* 70:137–139. <http://dx.doi.org/10.1111/j.1751-0813.1993.tb06105.x>.
 4. Sarker S, Patterson EI, Peters A, Baker GB, Forwood JK, Ghorashi SA, Holdsworth M, Baker R, Murray N, Raidal SR. 2014. Mutability dynamics of an emergent single stranded DNA virus in a naive host. *PLoS One* 9:e8537. <http://dx.doi.org/10.1371/journal.pone.0085370>.
 5. Sarker S, Ghorashi SA, Forwood JK, Raidal SR. 2014. Rapid genotyping of *beak and feather disease virus* using high-resolution DNA melt curve analysis. *J Virol Methods* 208:47–55. <http://dx.doi.org/10.1016/j.jviromet.2014.07.031>.
 6. Sarker S, Ghorashi SA, Forwood JK, Bent SJ, Peters A, Raidal SR. 2014. Phylogeny of *beak and feather disease virus* in cockatoos demonstrates host generalism and multiple-variant infections within *Psittaciformes*. *Virology* 460–461:72–88. <http://dx.doi.org/10.1016/j.virol.2014.04.021>.
 7. Sarker S, Lloyd C, Forwood J, Raidal SR. 2016. Forensic genetic evidence of beak and feather disease virus infection in a powerful owl (*Ninox strenua*). *Emu* 116:71–74. <http://dx.doi.org/10.1071/MU15063>.
 8. Das S, Sarker S, Peters A, Ghorashi SA, Phalen D, Forwood JK, Raidal SR. 2016. Evolution of circoviruses in lorikeets behind its hosts. *Mol Phylogenet Evol* 100:281–291. <http://dx.doi.org/10.1016/j.ympev.2016.04.024>.
 9. Sarker S, Forwood JK, Ghorashi SA, Peters A, Raidal SR. 2015. Beak and feather disease virus genotypes in Australian parrots reveals flexible host-switching. *Aust Vet J* 93:471–475. <http://dx.doi.org/10.1111/avj.12389>.
 10. Tracey JP, Woods R, Roshier D, West P, Saunders GR. 2004. 1004. The role of wild birds in the transmission of avian influenza for Australia: an ecological perspective. *Emu* 104:109–124. <http://dx.doi.org/10.1071/MU04017>.
 11. Bonne N, Clark P, Shearer P, Raidal S. 2008. Elimination of false-positive polymerase chain reaction results resulting from hole punch carryover contamination. *J Vet Diagn Invest* 20:60–63. <http://dx.doi.org/10.1177/104063870802000111>.
 12. Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2013. GenBank. *Nucleic Acids Res* 41:D36–D42. <http://dx.doi.org/10.1093/nar/gks1195>.
 13. Sarker S, Moylan KG, Ghorashi SA, Forwood JK, Peters A, Raidal SR. 2015. Evidence of a deep viral host switch event with *beak and feather disease virus* infection in rainbow bee-eaters (*Merops ornatus*). *Sci Rep* 5:14511. <http://dx.doi.org/10.1038/srep14511>.
 14. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
 15. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402. <http://dx.doi.org/10.1093/nar/25.17.3389>.