

## Allelopathic potential of pasture legume species and selected winter crops on barnyard grass and rice

Jhoana Opena<sup>1</sup>, Hanwen Wu<sup>1,2</sup>, Sergio Moroni<sup>1</sup>, Deirdre Lemerle<sup>1</sup> and James Pratley<sup>1</sup>

<sup>1</sup>Graham Centre for Agricultural Innovation (Charles Sturt University and NSW Department of Primary Industries), School of Agricultural and Wine Sciences, Locked Bag 588, Wagga Wagga, New South Wales 2678, Australia

<sup>2</sup>NSW Department of Primary Industries, Wagga Wagga 2650, New South Wales, Australia  
(jopena@csu.edu.au)

**Summary** Allelopathic potential of pasture legumes and selected winter crops on barnyard grass and rice were investigated. Three annual pasture legume species (antas subclover, shaftal persian, and viper balansa clovers), canola and barley were used for the evaluation of potential allelopathic effects of residues on the germination and early growth of barnyard grass and rice varieties used for drill sown rice such

as YRK5, YRM70, Opus and Sherpa. Results of the study will be presented during the conference. This information on the use of pasture legume species and/or winter crops in rotation with rice may provide a non-chemical alternative method to manage barnyard grass in drill-sown rice.

**Keywords** Allelopathy, barnyard grass, pasture legumes.