

International Journal of Research in Exercise Physiology

Original Research Article

The Effect of Aerobic Training Years on Systemic Oxygen Utilisation, and Peripheral O₂ Extraction in the *Vastus Lateralis*, *Gastrocnemius* and Pre-Frontal Cortex in Young Men

Gavin Buzza¹, Geoff P. Lovell², Christopher D. Askew³, Colin Solomon³

¹School of School of Exercise Science, Sport & Health, Charles Sturt University, Port Macquarie, Australia

²School of Social Sciences, University of the Sunshine Coast, Sippy Downs Australia

³School of Health and Sport Sciences, University of the Sunshine Coast, Sippy Downs, Australia

ABSTRACT

Introduction: This study was designed to determine if systemic O₂ utilisation ($\dot{V}O_2$), and tissue O₂ extraction (deoxyhemoglobin [HHb]) in the *vastus lateralis* (VL), *gastrocnemius* (GAST) and pre-frontal cortex (PFC) differed between aerobically short term trained (STT 6 - 24 months) and long term trained (LTT > 5 yr) men aged 18 - 30 years, who were matched for current training load.

Methods: Fourteen STT and 13 LTT participants completed ramp incremental (RI) and square-wave constant load (SWCL) (3 min at 25%, 80% and 25% and 20 min at 90% of ventilatory threshold [VT]) tests on a cycle ergometer. **Results:** ANCOVA revealed that $\dot{V}O_2$ was higher in LTT compared to STT in the RI ($p < 0.001$) and SWCL ($p = 0.004$) tests. There was no difference in Δ HHb in the VL, GAST or PFC between LTT and STT during both tests. However, there was a group x intensity interaction in the GAST during SWCL. **Conclusion:** These results suggest that in men aged 18 - 30 years, with up to 24 months of aerobic training, additional training years can increase systemic O₂ utilisation without increased current training load and that peripheral adaptations have little influence on the observed increases.

KEYWORDS: Aerobic training, oxygenation, $\dot{V}O_{2peak}$.

Introduction

Aerobic training consistently increases peak aerobic capacity ($\dot{V}O_{2peak}$) in healthy adults¹⁻³, with a dose response relationship existing (to a limit) between increased $\dot{V}O_{2peak}$ and the

training load (intensity, duration and frequency) of the intervention³⁻⁶. In previously untrained young men, training studies of between 9 - 52 weeks report a plateau in $\dot{V}O_{2peak}$ gains^{4,5,7}, however, this could be due to

the training stimulus not changing as adaptations occur. It is yet to be investigated if (independent of current training load) further gains in $\dot{V}O_{2peak}$ can result from additional training years in young men. If as suggested, the primary mediator of increases in $\dot{V}O_{2peak}$ is training load not years of training, the $\dot{V}O_{2peak}$ of young men that are matched for age and training load but differ in years of training should be similar. However, we expect that training years would affect $\dot{V}O_{2peak}$, as reported in similar studies of older⁸ and younger⁹ women from our laboratory

The gains in $\dot{V}O_{2peak}$ following aerobic training result from increased central O_2 delivery (cardiac output) to, and O_2 extraction (arterio-venous O_2 difference) by peripheral muscles. In young men, the relative contribution that central and peripheral adaptations make to increasing $\dot{V}O_{2peak}$ is not known and varies depending on the training load and duration of the intervention^{6,10-12}. Further, a meta-analysis of short term (5 - 13 wk) training studies reported that central adaptations are linearly associated with and likely responsible for increases in $\dot{V}O_{2peak}$, with little to no changes in peripheral O_2 extraction (arterio-venous O_2 difference)¹⁰. However, except for one study that assessed single limb O_2 uptake⁶, the other studies focused on changes in whole body O_2 extraction (arterio-venous O_2 difference) which reflects O_2 extraction of all tissues in the body, not isolated muscles.

Near-Infrared Spectroscopy (NIRS) provides a method of non-invasively monitoring changes in local tissue O_2 extraction, through changes in deoxyhemoglobin (HHb) and oxyhemoglobin (O_2Hb)¹³⁻¹⁵. The HHb pattern

reflects the dynamic balance between O_2 delivery and O_2 utilisation at the site of the measurement, and thus tissue O_2 extraction^{16,17}. Changes in HHb response (ΔHHb) has been used to report higher tissue O_2 extraction at the *vastus lateralis* (VL) of trained compared to untrained young and older men^{18,19} and young²⁰ and older²¹ women during transitions from low to moderate exercise, and to report a correlation between $\dot{V}O_{2peak}$ and peak HHb in the VL and *rectus femoris* muscles²². However, the effect of aerobic training years on peak muscle tissue O_2 extraction, or tissue O_2 extraction at a constant sub-ventilatory threshold (VT) load for an extended period remains to be investigated in a cohort of young men. Furthermore, the reported large decrease in O_2Hb and maintained or continued increases in HHb in the pre-frontal cortex (PFC) during high intensity exercise in highly trained ($\dot{V}O_{2peak}$ of $62 \pm 7.9 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$) but not untrained young men¹³ have led to theories of a potential relationship between PFC oxygenation and exercise performance^{14,23,24}. Therefore, a better understanding of tissue oxygenation responses at various sites during different intensity exercise, and following different durations (yr) of training may help clarify the influence of aerobic training on these determinants of aerobic capacity.

The aim of this current study was to simultaneously investigate the difference in $\dot{V}O_2$, and ΔHHb in the VL and *gastrocnemius* (GAST) muscles and the PFC between STT (6 - 24 months) and LTT (> 5 yr) aerobically trained young men aged 18 - 30 years who were matched for current training load. Participants completed ramp incremental (RI) and sub-VT square-wave constant load (SWCL) cycling

tests. It was hypothesised that: 1) $\dot{V}O_2$, and ΔHb would be higher in the VL and GAST of LTT compared to the STT during the RI and each measured relative intensity (25%, 80% and 90% VT) during SWCL; and 2), there would be no difference in ΔHb in the PFC between LTT and STT at any exercise intensity during both tests.

Methods

Ethical approval

This study was approved by the Human Research Ethics Committee at the University of the Sunshine Coast, Australia (S/14/676) and participants provided written informed consent.

Participants

The sample of young (18 - 30 yr) Caucasian men consisted of one group of 14 STT (6 - 24 months) men, and one group of 13 LTT (> 5 yr

men. Each participant reported having exceeded 200 min of moderate to vigorous aerobic training per week (including cycling), not missing more than two weeks of training over any six-month period, as determined from self-reported physical activity training logs (refer Table 1). Current training load (volume and exercise intensity) is a primary factor influencing $\dot{V}O_2$ and ΔHb during exercise, with both being higher in higher trained individuals^{18,22}. All participants reported their current training load as a typical training week across the previous six months. Participants' physical characteristics and training history are presented in Table 1. Following medical screening (Physical Activity Readiness Questionnaire²⁵ and Medical Health Questionnaire), exclusion included any health related issues or medications that would affect participant safety and or exercise capacity or O_2 utilisation/extraction.

Table 1. Participant characteristics for short term trained and long term trained young men.

Characteristic	STT	LTT
Age (yr)	23.5 (3.6)	24.0 (3.6)
Weight (kg)	84.3 (14.4)	79.7 (12.1)
Height (cm)	180.4 (8.8)	184.1 (8.7)
VL adipose (thickness mm)	7.4 (2.6)	3.9 (1.2) *
GAST adipose (thickness mm)	4.9 (3.0)	3.0 (0.6) *
Current training load (AU)	1189 (331)	1302 (324)
Current training (yr)	1.5 (0.5)	8.3 (2.7) *
VT $\dot{V}O_2$ (mL · kg ⁻¹ · min ⁻¹)	28.5 (4.7)	36.9 (5.2) *
VT % of Peak (mL · kg ⁻¹ · min ⁻¹)	62.8 (6.9)	65.5 (4.8)

Values are mean (SD). (STT n = 14, LTT n = 13)

* Significant ($p < 0.05$) difference between LTT and STT.

Current training (yr) = current number of continuous years of training; Current training load (AU) = volume x intensity (light = 1, moderate = 2 and high = 3).

Study design

The study used a cross-sectional, two group

(STT and LTT), within subject repeated measures (exercise intensity within RI and

SWCL) design. The independent variable used was current training duration (yr). The dependent variables were systemic $\dot{V}O_2$, heart rate (HR), rating of perceived exertion (RPE) (Borg's 1 - 10 category-ratio [CR-10])²⁶, and $\Delta H H b$ in the VL, GAST and PFC. The current training load was not significantly different between STT and LTT ($p = 0.909$).

Each participant completed two testing sessions in a temperature controlled (20 - 23° C) Exercise Physiology Laboratory. Prior to testing, participants were required to abstain from caffeine and food for four hours, and alcohol and intense exercise for 24 hours.

Procedures

Session One

The aim of session one was to determine participant characteristics, and $\dot{V}O_2$, HR, RPE (Table 2) and $\Delta H H b$ in the VL, GAST and PFC during a RI (increments = 1 W per 2-s) cycling test to volitional cessation on a Velotron cycle ergometer (Racermate, Seattle, USA). Anthropometric, pulmonary function (Spiro II spirometer, Medical International Research, Rome, Italy), HR and RPE data were recorded using standard measures as previously described⁸. The $\dot{V}O_2$, HR, and $\Delta H H b$ in the VL, GAST and PFC were recorded continuously during exercise, while RPE was recorded within the last 10 s of each minute (except for the warm-up) and at the point of volitional cessation. To encourage peak results, participants received verbal feedback and encouragement from the tester. Expired gas was analysed to determine $\dot{V}O_{2peak}$ and VT, with $\dot{V}O_{2peak}$ determined as the highest 15-s average $\dot{V}O_2$ value within the last minute of

this test, and VT determined using the V-slope method described by Beaver, Wasserman, Whipp²⁷. Briefly, this method defines the point where the rate of rise in CO_2 output ($\dot{V}CO_2$) relative to minute ventilation exceeds the relative rise in $\dot{V}O_2$. This VT was time aligned with the power (watts) of the cycle ergometer to calculate the power outputs for the subsequent SWCL test. As direct comparisons were not made between the RI and SWCL tests or for systemic $\dot{V}O_2$ and muscle O_2 extraction (i.e. $\dot{V}O_2$ to $\Delta H H b$ ratio), $\dot{V}O_{2peak}$ was not left shifted to accommodate for any potential phase I- phase II $\dot{V}O_2$ lag time.

Session Two

Session two was conducted three to 14 days after session one. The aim of session two was to determine the participants' $\dot{V}O_2$, HR, RPE (Table 3), and $\Delta H H b$ in the VL, GAST and PFC while cycling at a relative intensity of 25%, 80% and 90% of the VT power output obtained from the RI test. The timing and intensities for the SWCL test were; 3 minutes at 25%, 80%, 25%; 20 minutes at 90%; and a further 3 minutes at 25% of calculated VT.

The $\dot{V}O_2$, HR, and $\Delta H H b$ in the VL, GAST and PFC were recorded continuously during exercise, while RPE was recorded within the last 10 s of the third min of each of the three min SWCL stages, and every fourth min within the 20 min stage using methods as previously described⁸. Participants were not encouraged or provided feedback during the test, to minimise cognitive stimulus.

Tissue Deoxyhemoglobin

Tissue oxygenation (HHb and O_2Hb) data were measured simultaneously and continuously in

the left VL, GAST and PFC using a single-channel NIRS system (PortaMon and Portalite, Artinis Medical Systems BV, Zetten, Netherlands). The muscle optodes were fixed to the skin at the mid-belly of the muscle using adhesive tape and wrapped with low compression black elastic bandage, and the PFC optode was fixed to the skin at the left PFC using adhesive tape, then covered with a black headband⁸.

All NIRS primary data (ΔHHb and $\Delta\text{O}_2\text{Hb}$) were recorded at 10 Hz. The last 20-s of resting values were averaged to obtain baseline values. Any changes were then expressed relative to these baseline values and then calculated and displayed as follows: 15 s averages for RI; 30 s averages for SWCL; total average data for the 15-s preceding 90% VT and peak exercise for RI; and total data for each exercise intensity for SWCL. Compared to $\Delta\text{O}_2\text{Hb}$, ΔHHb (the primary variable for the present study) is less affected by changes in hemodynamics^{16,17,28} and is therefore a better indicator of tissue O_2 extraction. Therefore, only ΔHHb data have been presented in this present study.

Statistical Analyses

All statistical analyses were performed using SPSS (version 24, SPSS Inc., Chicago, IL). Prior to statistical analysis, data were checked for normality (Shapiro-Wilk test of normality and Mauchly test for Sphericity), and that the relevant assumptions for each test were met.

To identify the presence of any significant group (STT vs LTT) differences in current training load, independent t-tests were

conducted. To identify the presence of any significant group (STT vs LTT) and exercise intensity main effects or interactions, two-way Analysis of Covariance (ANCOVAs) were conducted on each dependent variable within each test. For the RI, 2 (group: STT and LTT) x 2 (intensity: 90% of VT and peak) ANCOVAs were performed, and for the SWCL, 2 (group: STT and LTT) x 3 (intensity: 25% [first bout at 25%] 80% and 90% of VT) ANCOVAs were performed. Due to the potential substantial effect of current training load on physiological responses to exercise, training load was included as a covariate in all analyses. The level for statistical significance was set to $p < 0.05$ for all analyses. Partial-eta squared was used to indicate the effect size as small ($\eta^2 > 0.01$), medium ($\eta^2 > 0.06$) or large ($\eta^2 > 0.14$) as per Cohen²⁹.

Results

The $\dot{V}\text{O}_{2\text{peak}}$, HR and RPE exhibited significant intensity main effects with large effect sizes in both the RI and SWCL tests, while the ΔHHb in the VL, GAST and PFC exhibited significant intensity main effects with large effect sizes in the SWCL but not the RI test. These variables increased with exercise intensity as physiologically expected (descriptive statistics are shown in Tables 2 and 3). Significant group main effects with large effect sizes were observed for $\dot{V}\text{O}_2$ in both the RI and SWCL tests, with $\dot{V}\text{O}_2$ higher in LTT. There were significant group by intensity interactions with large effect sizes for $\dot{V}\text{O}_2$, and ΔHHb in GAST in the SWCL test, with the difference increasing with increased intensity. Results for group main effects and group by intensity interactions are shown in Table 4.

Table 2. Systemic oxygen utilisation, ventilation, heart rate and rating of perceived exertion of short term trained and long term trained young men at 90% ventilatory threshold and peak exercise during Ramp Incremental cycling.

Variable	90% VT		Peak	
	STT	LTT	STT	LTT
$\dot{V}O_2$ (mL · kg ⁻¹ · min ⁻¹) ^{a, b}	26.0 (3.6)	33.8 (4.4)	45.2 (4.9)	56.3 (6.9)
HR (Beats · min ⁻¹) ^{#, b}	141.1 (13.7)	140.4 (11.4)	177.9 (9.8)	180.3 (10.3)
RPE ^b	4.2 (0.8)	4.5 (0.9)	9.9 (0.3)	9.8 (0.4)

Values are mean (SD).

^a significant group main effect; ^b significant intensity main effect; ^{ab} significant group by intensity interaction; * significant difference between STT and LTT $p < 0.05$; # HR at 90% VT, STT $n = 12$, LTT $n = 13$. HR at Peak, STT $n = 13$, LTT $n = 13$

VT: Ventilatory threshold; STT: Short term trained; LTT: Long term trained; $\dot{V}O_2$: Oxygen utilisation; HR: Heart rate; RPE: Rating of perceived exertion.

Table 3. Systemic oxygen utilisation, heart rate and rating of perceived exertion of short term trained and long term trained young men at 25%, 80% and 90% ventilatory threshold during Square-Wave Constant Load cycling.

Variable	25% VT		80% VT		90% VT	
	STT	LTT	STT	LTT	STT	LTT
$\dot{V}O_2$ (mL · kg ⁻¹ · min ⁻¹) ^{a, b, ab}	10.9 (1.6)	12.8 (2.2) *	23.0 (3.5)	28.2 (4.7) *	31.4 (5.6)	39.2 (7.4) *
HR (Beats · min ⁻¹) ^{#, b}	89.1 (8.5)	90 (7.3)	120.7 (10.8)	122.1 (7.6)	148.9 (14.5)	153.8 (12.5)
RPE ^b	1.1 (0.3)	1.2 (0.4)	3.5 (1.5)	3.2 (0.4)	5.8 (1.0)	5.8 (0.8)

Values are mean (SD).

^a significant group main effect; ^b significant intensity main effect; ^{ab} significant group by intensity interaction; * significant difference between STT and LTT $p < 0.05$; #HR at 25%, 80% and 90% VT; STT $n = 14$, LTT $n = 11$.

VT: Ventilatory threshold; STT: Short term trained; LTT: Long term trained; $\dot{V}O_2$: Oxygen utilisation; HR: Heart rate; RPE: Rating of perceived exertion.

Table 4. Results of two-way Analysis of Covariance (ANCOVAs) for systemic oxygen utilisation, heart rate, rating of perceived exertion and HHb of young men during ramp incremental and square-wave constant load cycling.

Variable	Test	Main effect for group					Group x intensity interaction				
		Df	F	p	η^2	β	Df	F	p	η^2	β
$\dot{V}O_2$ (mL · kg ⁻¹ · min ⁻¹)	RI	1, 24	24.412	< 0.001*	0.504	0.997	1, 24	2.952	0.099	0.110	0.378
	SWCL	1, 24	9.981	0.004*	0.294	0.858	2, 48	7.803	0.001*	0.245	0.939
HR (Beats · min ⁻¹)	RI	1, 22	0.005	0.994	< 0.001	0.051	1, 22	0.410	0.529	0.018	0.094
	SWCL	1, 22	0.435	0.516	0.019	0.097	2, 44	0.528	0.536	0.026	0.141
RPE	RI	1, 24	0.400	0.533	0.016	0.093	1, 24	1.702	0.204	0.066	0.240
	SWCL	1, 24	0.014	0.907	0.001	0.051	2, 48	0.298	0.744	0.012	0.095
VL HHb	RI	1, 23	0.450	0.509	0.019	0.099	1, 23	1.200	0.285	0.050	0.183
	SWCL	1, 24	0.230	0.636	0.009	0.075	2, 48	0.117	0.890	0.005	0.067
GAST HHb	RI	1, 22	3.899	0.061	0.151	0.471	1, 22	0.013	0.910	0.001	0.051
	SWCL	1, 22	2.284	0.145	0.094	0.304	2, 44	7.938	0.001*	0.265	0.941
PFC HHb	RI	1, 23	0.237	0.631	0.010	0.075	1, 23	0.145	0.707	0.006	0.065
	SWCL	1, 22	0.093	0.763	0.004	0.060	2, 44	0.066	0.936	0.003	0.059

*Significant p < 0.05

RI: Ramp incremental; SWCL: Square-wave constant load; $\dot{V}O_2$: Oxygen utilisation; HR: Heart rate; RPE: Rating of perceived exertion, VL: *Vastus lateralis*; GAST: *Gastrocnemius*; PFC: Pre-frontal cortex; HHb: deoxyhaemoglobin.




Figure 1. Mean (SD); Δ HHb in the VL, GAST and PFC during Ramp Incremental cycling. Panel (A) Δ HHb in the VL. Panel (B) Δ HHb in the GAST. Panel (C) Δ HHb in the PFC. STT 0 - 8.5 min (n = 13); 8.5 - 10.5 min (n = 7 - 11); 10.5 - 12.5 min (n = 3 - 6). LTT 0 - 9.25 min (n = 13); 9.25 - 11.75 (n = 9 - 12); 11.75 - 14 (n = 2 - 7). Square trace LTT, diamond trace STT.




Figure 2. Mean (SD); Δ HHb in the VL, GAST and PFC during Square-Wave Constant Load cycling. Panel (A) Δ HHb in the VL. Panel (B) Δ HHb in the GAST. Panel (C) Δ HHb in the PFC. Diamond shape STT, square shape LTT.




Figure 3. Mean (SD); Δ HHb in the VL, GAST and PFC during Ramp Incremental cycling. Panel (A) Δ HHb in the VL. Panel (B) Δ HHb in the GAST. Panel (C) Δ HHb in the PFC at 90% of VT and peak exercise. Pattern fill STT, solid fill LTT.




Figure 4. Mean (SD); Δ HHb in the VL, GAST and PFC during Square-Wave Constant Load cycling. Panel (A) Δ HHb in the VL. Panel (B) Δ HHb in the GAST. Panel (C) Δ HHb in the PFC at 90% of VT and peak exercise. Pattern fill STT, solid fill LTT.

* Significant ($p < 0.05$) differences between STT and LTT.

Discussion

This study examined the effect of aerobic training years (STT vs LTT) on $\dot{V}O_2$, and O_2 extraction (ΔHHb) in the VL, GAST and PFC at peak, and sub-VT exercise intensities. The primary findings were that: (i) $\dot{V}O_2$ was higher in LTT compared to STT in both the RI and SWCL tests; and (ii) although a significant group x intensity interaction was observed for ΔHHb in the GAST in the SWCL test, ΔHHb in the VL, GAST and PFC was not significantly different between LTT and STT in either test. This suggests that the higher $\dot{V}O_2$ in LTT was due to higher central O_2 delivery rather than peripheral O_2 extraction.

Systemic Oxygen Utilisation

As hypothesised, $\dot{V}O_{2peak}$ was significantly higher in LTT compared to STT. The $\dot{V}O_{2peak}$ values of $56.3 \pm 6.9 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ for LTT and $45.2 \pm 4.9 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ for STT are also consistent with the training load and experience reported by the participants³⁰. However, these present results do not support suggestions that in young men, $\dot{V}O_{2peak}$ plateaus within 12 months of commencing aerobic training⁴⁻⁵. While differences in current training years between previous studies and the present study provides a possible explanation for the difference in results, past training volumes of participants in the present study (which was not assessed) may have influenced the results. Additionally, compared to the STT, the genetic determinants of trainability of the LTT participants may have been stronger, and a reason why the LTT in the present study had continued aerobic training³¹⁻³².

Concomitant to $\dot{V}O_{2peak}$, as expected, $\dot{V}O_2$ at VT was higher in the LTT compared to the STT. However, there was no difference when the $\dot{V}O_2$ was expressed as a percentage of $\dot{V}O_{2peak}$, indicating increases in VT paralleled that of $\dot{V}O_{2peak}$. As some training methods such as high intensity interval training increase VT as a percentage of $\dot{V}O_{2peak}$ ³³⁻³⁴, this finding may suggest that the training stimulus of both LTT and STT were similar.

Despite exercising at the same calculated relative exercise intensity during the study, $\dot{V}O_2$ was higher in LTT compared to STT during the SWCL test. This difference increased with increased exercise intensity from $1.9 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ at 25% to $5.2 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ at 80% and $11.1 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ at 90% of VT. This combined with other findings in older men¹⁸ and older^{8,21} and younger women⁹ suggests that regardless of age and sex, aerobic training increases systemic O_2 utilisation while exercising at constant loads below VT.

In the present study, heart rates were not different between LTT and STT at any intensity in either test, supporting that LTT and STT were matched for age and were cycling at similar relative intensities during the SWCL test. Although peak heart rates are a function of age and not training, training induced faster heart rate dynamics during transitions from lower to higher exercise intensities could impact the delivery of substrates or O_2 required for oxidative phosphorylation^{3,18}. Additionally, RPE was not different at any intensity during the SWCL test, further supporting that exercise intensities were matched between STT and LTT.

Tissue Deoxyhemoglobin

For ΔHHb in the VL and GAST, the present data were contrary to the hypothesis with ΔHHb not being different between LTT and STT during either test (Figures 2 and 3). This suggests that in young men, beyond approximately two years of aerobic training, additional training years does not independently further increase O_2 extraction (ΔHHb) in the VL or GAST. It has been suggested that muscle tissue oxygenation (ΔHHb and $\Delta\text{O}_2\text{Hb}$) correlates with $\dot{V}\text{O}_{2\text{peak}}$ in young men^{22,35}. However, the results of the present study do not support this, with ΔHHb in the VL and GAST being not different between LTT and STT at peak exercise despite the higher $\dot{V}\text{O}_2$ in the LLT.

The effect of exercise training on the peak HHb in the VL of young men has previously only been reported in one study in untrained young men. Following a short term (6 wk) high intensity running program, peak HHb was significantly higher during cycling exercise³³. In addition, changes in sub-maximal HHb have been limited to studies comparing changes in the ratio of tau (response time to a step increment in work load) HHb in the VL to tau pulmonary $\dot{V}\text{O}_2$ (tau ΔHHb :tau $\Delta\dot{V}\text{O}_2$) during transitions from low to moderate intensity exercise^{18-19,36-37}. The reduced $\Delta\text{HHb}:\Delta\dot{V}\text{O}_2$ ratio was reported to reflect an improved matching of O_2 distribution to extraction. However, tau HHb was unchanged in these previous studies^{18-19,36-37}.

In the present study, the difference in ΔHHb in the GAST between STT and LTT did not reach statistical significance. However, the ΔHHb was substantially higher in LTT at 90% VT and

peak exercise during RI, and at 80% and 90% during SWCL, and therefore, likely to be physiologically meaningful. Due to the high between-participant variability in ΔHHb during testing, it is possible that the results may not be completely indicative of potential differences. Three studies have previously published oxygenation (HHb or O_2Hb) changes in the GAST during exercise^{8,38-39}. However, only one study from our laboratory investigated the effect of aerobic training⁸. That study showed no significant difference in ΔHHb in the GAST between STT and LTT older women at peak or sub-VT exercise. This, in addition to studies in our laboratory of older men and young women showing differences in ΔHHb in the GAST between LTT compared to STT suggests that training induced changes in O_2 extraction in the GAST may differ as a function of age, sex and exercise intensity. Possible explanations for these varied results include sex differences in muscle fibre distribution⁴⁰, and age-related reduction in leg blood flow and perfusion pressure in older women⁴¹.

In the present study, the ΔHHb pattern in the GAST at the onset of exercise opposed that of the VL. That is, ΔHHb in the VL gradually increasing from the onset of exercise to a plateau, whereas ΔHHb in the GAST decreased substantially at the onset of exercise then progressively increased with increased exercise intensity (Figures 1 and 2). This response has previously been reported in older women⁸ and young women⁹. As metabolic and muscle fibre recruitment processes are influenced by fibre type characteristics (slow-twitch, low O_2 extraction to fast-twitch, higher O_2 extraction), variations

in muscle fibre characteristics between the VL and GAST may provide a possible explanation for the differences in HHb patterns^{13,40,42-43}.

For HHb in the PFC, the present results support the hypothesis in that ΔHHb was not different between STT and LTT in either test. This supports recent consensus that cerebral oxygenation does not limit the exercise performance of healthy young adults under normoxic conditions^{13,44}. A link between cerebral oxygenation, central motor drive and exercise performance emerged following reports of reduced O_2Hb and increased HHb near maximal exercise^{14,23-24,45-48}, and increased performance following administration of O_2 near maximal exercise in hypoxic^{23,49} and normoxic conditions⁵⁰. The present results oppose suggestions that training status (aerobic capacity) impacts cerebral oxygenation^{48,51}. Indeed, our data showing $\dot{V}\text{O}_2$ values differed between groups and ΔHHb in the PFC did not differ, suggests that aerobic capacity does not affect PFC oxygenation. However, it is possible that the difference in aerobic capacity between LTT and STT was not great enough to impact PFC oxygenation. Given that aerobic capacity only appears to impact cerebral oxygenation in high level athletes¹³, it is also possible that true maximum effort was not achieved, and/or aerobic capacity was not high enough to impact PFC oxygenation.

LIMITATIONS

A potential limitation of this present study is that while regular cycling was a requirement for recruitment, the contribution cycling had on individual training volumes was not determined. Furthermore, the adipose tissue

thickness of the STT was higher than the LTT. However, while high levels of adipose tissue may affect NIRS measures, all participants were within the recommended ranges (< 34 mm) for ΔHHb not to be affected⁵².

PRACTICAL IMPLICATIONS AND FUTURE RESEARCH

The results of this study have implications for aerobic training in young men. Specifically, the present data suggest additional years of regular training in young men, without increasing weekly training load, may provide significant further increases in $\dot{V}\text{O}_2$, with the increase resulting predominately from central not peripheral adaptations. Moreover, $\Delta\text{PFC O}_2$ extraction does not change with additional years of training in young men aged 18 – 30 yr. Future research should include training studies that exceed 12 months duration and monitor central delivery to and extraction of O_2 in multiple peripheral tissues during various absolute and relative exercise intensities.

Conclusions

It is concluded that independent of current training load, regular aerobic training beyond approximately two years duration significantly increases the $\dot{V}\text{O}_2$ of young men during ramp incremental and square-wave constant load exercise. This increase occurs without increases in tissue O_2 extraction in the *vastus lateralis* and *gastrocnemius* muscles.

Acknowledgements

Hugo Kerhervé and Yuri Kriel provided technical assistance.

Competing interests

This work did not receive financial support or funding.

Address for Correspondence

Gavin Buzza PhD. School of Exercise Science, Sport & Health, Charles Sturt University, Port Macquarie, NSW 2444, Australia. E-mail: gbuzza@csu.edu.au

References

1. Montero D, Diaz-Cañestro C. Endurance training and maximal oxygen consumption with ageing: Role of maximal cardiac output and oxygen extraction. *Eur J Prev Cardiol.* 2016;23(7):733-743.
2. Bouchard C, Rankinen T. Individual differences in response to regular physical activity. *Med Sci Sports Exerc.* 2001;33(6 Suppl):S446-51; discussion S52-S53.
3. Murias JM, Kowalchuk JM, Paterson DH. Time course and mechanisms of adaptations in cardiorespiratory fitness with endurance training in older and young men. *J Appl Physiol.* 2010;108(3):621-627.
4. Gaesser GA, Rich RG. Effects of high- and low-intensity exercise training on aerobic capacity and blood lipids. *Med Sci Sports Exerc.* 1984;16(3):269-274.
5. Arbab-Zadeh A, Perhonen M, Howden E, Peshock RM, Zhang R, Adams-Huet B, et al. Cardiac remodeling in response to 1 year of intensive endurance training. *Circulation.* 2014;130(24):2152-2161.
6. Roca J, Agusti AG, Alonso A, Poole DC, Viegas C, Barbera JA, et al. Effects of training on muscle O₂ transport at VO₂max. *J Appl Physiol.* 1992;73(3):1067-1076.
7. Howden EJ, Perhonen M, Peshock RM, Zhang R, Arbab-Zadeh A, Adams-Huet B, et al. Females have a blunted cardiovascular response to one year of intensive supervised endurance training. *J Appl Physiol.* 2015;119(1):37-46.
8. Buzza G, Lovell GP, Askew CD, Kerherve H, Solomon C. The effect of short and long term endurance training on systemic, and muscle and prefrontal cortex tissue oxygen utilisation in 40 - 60 year old women. *PLoS One.* 2016;11(11):e0165433.
9. Buzza G, Lovell GP, Askew CD, Solomon C. The Effect of Short- and Long-Term Aerobic Training Years on Systemic O₂ Utilization, and Muscle and Prefrontal Cortex Tissue Oxygen Extraction in Young Women. *J Strength Cond Res.* 2019;33(8):2128-2137.
10. Montero D, Diaz-Cañestro C, Lundby C. Endurance training and VO₂max: Role of maximal cardiac output and oxygen extraction. *Med Sci Sports Exerc.* 2015;47(10):2024-2033.
11. Wang E, Naess MS, Hoff J, Albert TL, Pham Q, Richardson RS, et al. Exercise-training-induced changes in metabolic capacity with age: the role of central cardiovascular plasticity. *Age.* 2014;36(2):665-676.
12. Ogawa T, Spina RJ, Martin WH, 3rd, Kohrt WM, Schechtman KB, Holloszy JO, et al. Effects of aging, sex, and physical training on cardiovascular responses to exercise. *Circulation.* 1992;86(2):494-503.
13. Boone J, Vandekerckhove K, Coomans I, Prieur F, Bourgois JG. An integrated view on the oxygenation responses to incremental exercise at the brain, the locomotor and respiratory muscles. *Eur J Appl Physiol.* 2016;116(11-12):2085-2102.
14. Bhambhani Y, Malik R, Mookerjee S. Cerebral oxygenation declines at exercise intensities above the respiratory compensation threshold. *Resp Physiol Neurobiol.* 2007;156(2):196-202.
15. Ferrari M, Mottola L, Quaresima V. Principles, techniques, and limitations of near infrared spectroscopy. *Can J Appl Physiol.* 2004;29(4):463-487.
16. Grassi B, Pogliaghi S, Rampichini S, Quaresima V, Ferrari M, Marconi C, et al. Muscle oxygenation and pulmonary gas exchange kinetics during cycling exercise on-transitions in humans. *J Appl Physiol.* 2003;95(1):149-158.
17. DeLorey DS, Kowalchuk JM, Paterson DH. Relationship between pulmonary O₂ uptake kinetics and muscle deoxygenation during moderate-intensity exercise. *J Appl Physiol.* 2003;95(1):113-120.
18. Grey TM, Spencer MD, Belfry GR, Kowalchuk JM, Paterson DH, Murias JM. Effects of age and long-term endurance training on VO₂ kinetics. *Med Sci Sports Exerc.* 2015;47(2):289-298.
19. Murias JM, Kowalchuk JM, Paterson DH. Speeding of VO₂ kinetics with endurance training in old and young men is associated with improved matching of local O₂ delivery to muscle O₂ utilization. *J Appl Physiol.* 2010;108(4):913-922.
20. Murias JM, Kowalchuk JM, Paterson DH. Speeding of VO₂ kinetics in response to endurance-training in older and young women. *Eur J Appl Physiol.* 2011;111(2):235-243.
21. Dogra S, Spencer MD, Murias JM, Paterson DH. Oxygen uptake kinetics in endurance-trained and untrained postmenopausal women. *Appl Physiol Nutr Metab.* 2013;38(2):154-160.

22. Okushima D, Poole DC, Barstow TJ, Rossiter HB, Kondo N, Bowen TS, et al. Greater VO_2 peak is correlated with greater skeletal muscle deoxygenation amplitude and hemoglobin concentration within individual muscles during ramp-incremental cycle exercise. *Physiol Rep.* 2016;4(23):e13065.
23. Subudhi AW, Lorenz MC, Fulco CS, Roach RC. Cerebrovascular responses to incremental exercise during hypobaric hypoxia: effect of oxygenation on maximal performance. *Am J Physiol-Heart Circul Physiol.* 2008;294(1):H164-H171.
24. Rupp T, Perrey S. Prefrontal cortex oxygenation and neuromuscular responses to exhaustive exercise. *Eur J Appl Physiol.* 2008;102(2):153-163.
25. Bredin SSD, Gledhill N, Jamnik VK, Warburton DER. New risk stratification and physical activity clearance strategy for physicians and patients alike. *Can Fam Physician.* 2013;59(3):273-277.
26. Borg GA. Perceived exertion. *Exerc Sport Sci Rev.* 1974;2:131-153.
27. Beaver WL, Wasserman K, Whipp BJ. A new method for detecting anaerobic threshold by gas exchange. *J Appl Physiol.* 1986;60(6):2020-2027.
28. Ferreira LF, Townsend DK, Lutjemeier BJ, Barstow TJ. Muscle capillary blood flow kinetics estimated from pulmonary O_2 uptake and near-infrared spectroscopy. *J Appl Physiol.* 2005;98(5):1820-1828.
29. Cohen J. *Statistical power analysis for the behavioral sciences.* 2nd ed: Hillsdale, N.J. : L. Erlbaum Associates; 1988.
30. Edvardsen E, Scient C, Hansen BH, Holme IM, Dyrstad SM, Anderssen SA. Reference values for cardiorespiratory response and fitness on the treadmill in a 20- to 85-year-old population. *Chest.* 2013;144(1):241-248.
31. Bouchard C, Sarzynski MA, Rice TK, Kraus WE, Church TS, Sung YJ, et al. Genomic predictors of the maximal O_2 uptake response to standardized exercise training programs. *J Appl Physiol.* 2011;110(5):1160-1170.
32. Bacon AP, Carter RE, Ogle EA, Joyner MJ. VO_2 max trainability and high intensity interval training in humans: a meta-analysis. *PLoS One.* 2013;8(9):e73182.
33. Prieur F, Mucci P. Effect of high-intensity interval training on the profile of muscle deoxygenation heterogeneity during incremental exercise. *Eur J Appl Physiol.* 2013;113(1):249-257.
34. Sheykhlovand M, Gharaat M, Khalili E, Agha-Alinejad H. The effect of high-intensity interval training on ventilatory threshold and aerobic power in well-trained canoe polo athletes. *Sci Sports.* 2016;31(5):283-289.
35. Boone J, Barstow TJ, Celie B, Prieur F, Bourgois J. The interrelationship between muscle oxygenation, muscle activation, and pulmonary oxygen uptake to incremental ramp exercise: influence of aerobic fitness. *Appl Physiol Nutr Metab.* 2016;41(1):55-62.
36. Williams AM, Paterson DH, Kowalchuk JM. High-intensity interval training speeds the adjustment of pulmonary O_2 uptake, but not muscle deoxygenation, during moderate-intensity exercise transitions initiated from low and elevated baseline metabolic rates. *J Appl Physiol.* 2013;114(11):1550-1162.
37. McKay BR, Paterson DH, Kowalchuk JM. Effect of short-term high-intensity interval training vs. continuous training on O_2 uptake kinetics, muscle deoxygenation, and exercise performance. *J Appl Physiol.* 2009;107(1):128-138.
38. Hiroyuki H, Hamaoka T, Sako T, Nishio S, Kime R, Murakami M, et al. Oxygenation in vastus lateralis and lateral head of gastrocnemius during treadmill walking and running in humans. *Eur J Appl Physiol.* 2002;87(4-5):343-349.
39. Kriel Y, Kerherve HA, Askew CD, Solomon C. The effect of active versus passive recovery periods during high intensity intermittent exercise on local tissue oxygenation in 18 - 30 year old sedentary men. *PLoS One.* 2016;11(9):e0163733.
40. Hunter GR, Bamman MM, Larson-Meyer DE, Joanisse DR, McCarthy JP, Blaudeau TE, et al. Inverse relationship between exercise economy and oxidative capacity in muscle. *Eur J Appl Physiol.* 2005;94(5-6):558-568.
41. Proctor DN, Koch DW, Newcomer SC, Le KU, Smithmyer SL, Leuenberger UA. Leg blood flow and VO_2 during peak cycle exercise in younger and older women. *Med Sci Sports Exerc.* 2004;36(4):623-631.
42. Houmard JA, Weidner ML, Gavigan KE, Tyndall GL, Hickey MS, Alshami A. Fiber type and citrate synthase activity in the human gastrocnemius and vastus lateralis with aging. *J Appl Physiol.* 1998;85(4):1337-1341.
43. Edgerton VR, Smith JL, Simpson DR. Muscle fibre type populations of human leg muscles. *Histochem J.* 1975;7(3):259-266.
44. Subudhi AW, Olin JT, Dimmen AC, Polaner DM, Kayser B, Roach RC. Does cerebral oxygen delivery limit incremental exercise performance? *J Appl Physiol.* 2011;111(6):1727-1734.

45. Rasmussen P, Stie H, Nielsen B, Nybo L. Enhanced cerebral CO₂ reactivity during strenuous exercise in man. *Eur J Appl Physiol.* 2006;96(3):299-304.
46. Subudhi AW, Dimmen AC, Roach RC. Effects of acute hypoxia on cerebral and muscle oxygenation during incremental exercise. *J Appl Physiol.* 2007;103(1):177-183.
47. Smith KJ, Billaut F. Influence of cerebral and muscle oxygenation on repeated-sprint ability. *Eur J Appl Physiol.* 2010;109(5):989-999.
48. Rooks CR, Thom NJ, McCully KK, Dishman RK. Effects of incremental exercise on cerebral oxygenation measured by near-infrared spectroscopy: a systematic review. *Prog Neurobiol.* 2010;92(2):134-150.
49. Amann M, Calbet JAL. Convective oxygen transport and fatigue. *J Appl Physiol.* 2008;104(3):861-870.
50. Oussaidene K, Prieur F, Bougault V, Borel B, Matran R, Mucci P. Cerebral oxygenation during hyperoxia-induced increase in exercise tolerance for untrained men. *Eur J Appl Physiol.* 2013;113(8):2047-2056.
51. Oussaidene K, Tagougui S, Abaidia A, Matran R, Mucci P, Prieur F. Aerobic fitness influences cerebral oxygenation response to maximal exercise in healthy subjects. *Respir Physiol Neurobiol.* 2015;205:53-60.
52. van Beekvelt MC, Borghuis MS, van Engelen BG, Wevers RA, Colier WN. Adipose tissue thickness affects in vivo quantitative near-IR spectroscopy in human skeletal muscle. *Clin Sci.* 2001;101(1):21-28.