

This article is downloaded from



<http://researchoutput.csu.edu.au>

It is the paper published as:

Author: C. Gaden, J. M. Scott, K. Behrendt, A. Colvin, F. Scott, L. Shakhane, C. Guppy and J. Hoad

Title: Integrated overview of results from a farmlet experiment which compared the effects of pasture inputs and grazing management on profitability and sustainability

Journal: Animal Production Science

ISSN: 1836-0939 1836-5787

Year: 2013

Volume: 53

Pages: 841-855

Abstract: This paper provides an integrated overview of the results collated from component papers and discusses the inferences which can be drawn from what was a complex, agro ecosystem trial. The measurements recorded both early and late in the trial were tabulated for each of the farmlets and compared to each other as relative proportions, allowing visual presentation on a common, indexed scale. Because of equivalent starting conditions, there was little difference between farmlets early in the trial period (2000-2001) across a wide array of measured parameters including herbage mass, potential pasture growth rate, live weight, wool production per head, stocking rate, gross margin and equity. Although the trial experienced drier-than-average conditions, marked differences emerged between farmlets over time, due to the effects of treatments. During the latter half of the trial period (2003-2006), farmlet A showed numerous positive and a few negative consequences of the higher rate of pasture renovation and increased soil fertility compared to the other two farmlets. Whilst intensive rotational grazing resulted in superior control of gastrointestinal nematodes and slightly finer wool, this system had few effects on pastures and no positive effects on sheep live weights, wool production or stocking rate. Whereas farmlet A showed higher gross margins, it had a negative and much lower short-term cash position compared to farmlets B and C, due largely to the artificially high rate of pasture renovation undertaken on this farmlet during the trial. Although farmlet B had the highest cash position at the end of the trial, this came at a cost of the declining quality of its pastures. Modelling of the farmlet systems allowed the results of this drier-than-average experimental period to be seen in the context of long-term climatic expectations. The main factors responsible for lifting the productivity of farmlet A were the sowing of temperate species and increased soil fertility which enhanced the amount of legume and increased pasture quality and potential pasture growth. The factor which affected farmlet C most was the low proportion of the farmlet grazed at any one time, with high stock density imposed during grazing, which decreased feed intake quality. The paper concludes that more profitable and sustainable outcomes are most likely to arise from grazing enterprises which are proactively managed towards optimal outcomes by maintaining sufficient desirable perennial grasses with adequate legume content, enhancing soil fertility and employing flexible rotational grazing.

DOI/URL: <http://dx.doi.org/10.1071/AN12284> http://researchoutput.csu.edu.au/R/-?func=dbin-jump-full&object_id=46718&local_base=GEN01-CSU01

Author Address: kbehrendt@csu.edu.au

CRO Number: 46718

1 **Integrated overview of results from a farmlet experiment which**
2 **compared the effects of pasture inputs and grazing management on**
3 **profitability and sustainability**

4 JM Scott^{AN}, K Behrendt^B, A Colvin^C, F Scott^D, LM Shakhane^E, C Guppy^A, J Hoad^A, CA Gaden^F,
5 C Edwards^G, GN Hinch^A, OJ Cacho^H, GE Donald^J, D Cottle^A, T Coventry^K, G Williams^L, DF
6 Mackay^M

7 ^A School of Environmental and Rural Science, University of New England, Armidale, 2351, Australia

8 ^B EH Graham Centre for Agricultural Innovation (NSW Department of Primary Industries and Charles Sturt University),
9 School of Agricultural and Wine Science, Charles Sturt University, Orange, NSW 2800, Australia

10 ^C "Pine Grove" 6248 New England Highway, Ben Lomond NSW 2365, Australia

11 ^D NSW Trade & Investment, 4 Marsden Park Road, Calala NSW 2340, Australia

12 ^E National University of Lesotho, Lesotho

13 ^F "Beaumont", Invergowrie, NSW, 2350, Australia

14 ^G NSW Department of Primary Industries, Armidale, NSW 2350, Australia

15 ^H UNE Business School, University of New England, Armidale, 2351, Australia

16 ^J CSIRO Livestock Industries, Armidale, 2350, Australia

17 ^K Chairman, Cicerone Project, "Bailey Park", Armidale, NSW, 2350, Australia

18 ^L "Eastlake", Uralla, NSW, 2358, Australia

19 ^M 3 Jayne Cl, Armidale, NSW 2350, Australia

20 ^N Corresponding author: Jim.Scott@une.edu.au

21 **Abstract**

22 The Cicerone Project conducted a grazed farmlet experiment on the Northern Tablelands of
23 NSW, Australia, from July 2000 to December 2006 to address questions raised by local graziers
24 concerning how they might improve the profitability and sustainability of their grazing enterprises.
25 This unreplicated experiment examined three management systems at a whole farmlet scale: the
26 control (farmlet B) represented typical management for the region with flexible rotational grazing
27 and moderate inputs. A second (farmlet A) also used flexible rotational grazing but had a higher
28 level of pasture renovation and soil fertility, whilst the third (farmlet C) had the same moderate
29 inputs as farmlet B but employed intensive rotational grazing.

30 This paper provides an integrated overview of the results collated from component papers and
31 discusses the inferences which can be drawn from what was a complex, agroecosystem trial. The
32 measurements recorded both early and late in the trial were tabulated for each of the farmlets and
33 compared to each other as relative proportions, allowing visual presentation on a common, indexed
34 scale.

35 Because of equivalent starting conditions, there was little difference between farmlets early in
36 the trial period (2000-2001) across a wide array of measured parameters including herbage mass,
37 potential pasture growth rate, liveweight, wool production per head, stocking rate, gross margin
38 and equity. Although the trial experienced drier-than-average conditions, marked differences
39 emerged between farmlets over time, due to the effects of treatments. During the latter half of the
40 trial period (2003-2006), farmlet A showed numerous positive and a few negative consequences of
41 the higher rate of pasture renovation and increased soil fertility compared to the other two farmlets.
42 Whilst intensive rotational grazing resulted in superior control of gastrointestinal nematodes and
43 slightly finer wool, this system had few effects on pastures and no positive effects on sheep
44 liveweights, wool production or stocking rate.

45 Whereas farmlet A showed higher gross margins, it had a negative and much lower short-term
46 cash position compared to farmlets B and C, due largely to the artificially high rate of pasture
47 renovation undertaken on this farmlet during the trial. Although farmlet B had the highest cash
48 position at the end of the trial, this came at a cost of the declining quality of its pastures.
49 Modelling of the farmlet systems allowed the results of this drier-than-average experimental period
50 to be seen in the context of long-term climatic expectations.

51 The main factors responsible for lifting the productivity of farmlet A were the sowing of
52 temperate species and increased soil fertility which enhanced the amount of legume and increased
53 pasture quality and potential pasture growth. The factor which affected farmlet C most was the
54 low proportion of the farmlet grazed at any one time, with high stock density imposed during
55 grazing, which decreased feed intake quality.

56 The paper concludes that more profitable and sustainable outcomes are most likely to arise from
57 grazing enterprises which are proactively managed towards optimal outcomes by maintaining
58 sufficient desirable perennial grasses with adequate legume content, enhancing soil fertility and
59 employing flexible rotational grazing.

60 **Key Words**

61 Farming systems, multi-disciplinary, pasture quality, pasture legumes, parasitology, modelling,
62 risk, optimisation.

63 **Introduction**

64 This paper aims to integrate the findings from a series of related papers which were part of a
65 multi-disciplinary study of different grazed farmlets conducted on the Northern Tablelands of
66 NSW, Australia from July 2000 to December 2006. The experiment was set up to answer
67 questions chosen by local livestock producers about ways of enhancing the feed supply, either
68 through pasture renovation and soil fertility or through intensive grazing management.

69 Managing a grazing enterprise is challenging under any circumstances and especially so when
70 climatic conditions are highly variable, as they are in this region (Behrendt *et al.* 2012c; Sutherland
71 *et al.* 2012). According to Williams (1994), graziers have to learn to ‘simultaneously balance
72 many balls in the air’ in order to satisfactorily manage complex grazing systems. Lodge *et al.*
73 (1998) described some of the paddock management factors that are readily observed by farmers as:
74 the proportions of desirable perennial grasses and legumes, the amounts of green leaf and surface
75 litter, and the level of ground cover. To this list could be added observable attributes of livestock
76 such as condition, liveweight, pregnancy and general animal health. However, when one considers
77 additional influences on a farm system such as soil fertility, pasture composition, grazing
78 management and the stochastic (random) behaviour of prices and climate, it is clear that making
79 optimal decisions presents an intractable challenge for most managers.

80 In this paper we have summarised the relative contributions of soil, pasture, animal, economic
81 and environmental parameters over time in a similar fashion to the measurements of sustainability
82 reported by both Scott *et al.* (2000) and Lodge *et al.* (2003). This approach to quantifying

83 sustainability has been found to be useful not only for scientists but also for graziers whose visual
84 assessments of pastures have been found to be highly correlated with research assessments (Lodge
85 2002).

86 The main conclusion from the recent national Sustainable Grazing Systems experiment,
87 conducted across southern Australia, was that the productivity and sustainability of pasture-based
88 systems can be enhanced by higher levels of soil fertility, the amelioration of low soil pH, the
89 sowing of deep-rooted perennial grasses and the use of grazing methods that permit substantial rest
90 periods between grazings (Andrew *et al.* 2003). On the Northern Tablelands of NSW, there has
91 been a considerable body of research into the growth of both sown and native pastures and their
92 responses to nutrients (Cook *et al.* 1976; 1978; Lazenby and Lovett 1975; Robinson and Lazenby
93 1976; Whalley *et al.* 1976; Wolfe and Lazenby 1973). In spite of this published evidence that
94 substantial responses can be gained from the sowing of pastures and the amelioration of nutrient
95 deficiencies, many graziers in this region today question the long-term economic benefits of these
96 technologies, given the perceived high costs of improving pastures (Vere and Campbell 2004),
97 whilst many have expressed interest in intensive grazing management as a potential alternative
98 management solution (Scott *et al.* 2012c).

99 Comprehensive multi-disciplinary studies of grazing management are rare in Australia. One of
100 the earliest and most complete studies was a replicated trial conducted by Moore *et al.* (1946)
101 which compared continuous and rotational grazing over either 4- or 8-week intervals. They found
102 that, apart from retaining a somewhat higher level of lucerne in the pasture, rotational grazing was
103 not a reliable way to increase livestock production. However, as noted by Hacker (1993), there has
104 also been relatively little research into *intensive* rotational grazing (IRG) systems in Australia.
105 Also, most of the studies undertaken have compared such systems with *continuous* grazing
106 (Dowling *et al.* 2005; Earl and Jones 1996; Sanjari *et al.* 2008; Waller *et al.* 2001). Livestock
107 producer members of the Cicerone Project felt that, at least for the Northern Tablelands region of
108 NSW, which is part of Australia's temperate high rainfall zone, the use of *continuous* grazing as

109 the control treatment is inappropriate, as few graziers practise ‘continuous stocking’ as
110 implemented in these earlier experiments.

111 Whilst there are recorded cases of farm managers and proponents of intensive grazing
112 management claiming benefits of such systems including increased soil phosphorus, stocking rates
113 and profits (Cawood 2004; McCosker 2000), there are also publications where some of the claims
114 have been refuted (Dowling *et al.* 2005; Hall *et al.* 2011; Waugh 1997). The earliest of these
115 papers, written by a grazier from central NSW, reported unsatisfactory cattle growth when he
116 implemented ‘time control grazing’ on his property. After this experience, he resorted to a less
117 intensive form of rotational grazing with a rest period of about 30 days (Waugh 1997). In a recent
118 report of grazing systems on rangelands across inland Queensland, Hall *et al.* (2011) found that
119 stocking rate was a much more important driver of performance than grazing system. However,
120 shortly after the release of these findings, the methodology used in the project was reportedly
121 criticised by proponents of time control grazing and holistic resource management (Cawood 2011).

122 The debate about rotational grazing remains unresolved, partly because research has not
123 adequately addressed the human variables which affect management systems (Briske *et al.* 2011).
124 However, those planning the Cicerone farmllet experiment did consider the decision-making
125 approaches of livestock producers in designing the farmllet treatments (Scott *et al.* 2012c). During
126 the planning phase, Cicerone producer members and collaborators agreed that the research
127 conducted into intensive rotational grazing systems in the high rainfall zone of Australia had not
128 been conducted in a sufficiently comprehensive fashion to satisfactorily answer the questions that
129 members had. Whilst Norton (1998), in his extensive review of the literature, hypothesised that
130 intensive grazing management might result in increased production of pasture which could lead to
131 increased stocking rates, the Cicerone Project decided that such suggestions needed to be tested
132 under realistic experimental conditions. Thus, livestock producers requested that both pasture
133 renovation with higher soil fertility and intensive rotational grazing be compared with a more
134 typical management system within a whole farmllet experiment. This would permit the multiple
135 facets of such systems to be measured at a credible scale (Scott *et al.* 2012c).

136 The Cicerone farmlet experiment can be seen as an agroecological experiment with interactions
137 between the many component parts of grazed farming systems interacting in complex ways with
138 the climate and management. Eberhardt and Thomas (1991) pointed out that the design of
139 adequate ecosystem experiments often means that conventional experimental design criteria need
140 to be challenged; they also highlighted the need for more care to be taken in drawing inference
141 about cause and effect from such experiments due to the many complex interactions within
142 agricultural ecosystems. Details of the selection of experimental treatments and hypotheses
143 employed in this experiment have been described in detail by Scott *et al.* (2012c). In brief, the
144 hypothesis of the Cicerone farmlet experiment was that, compared to the typical farmlet (B), higher
145 pasture inputs combined with higher soil fertility (farmlet A) and/or intensive grazing management
146 (farmlet C) will result in a more profitable and sustainable enterprise.

147 **Methods**

148 The general methods adapted for use in the Cicerone farmlet trial have been described by Scott
149 *et al.* (2012c) whereas other more specific methods are contained in related component papers in
150 this Special Issue. The approach taken here has been to summarise the results and calculate indices
151 for all measured parameters so that readers can assess for themselves the validity of the
152 interpretations of evidence drawn by the authors of this paper.

153 This method of comparing treatments across a range of criteria is based on an approach
154 developed for an earlier grazed trial which aimed to quantify sustainability (Scott *et al.* 2000).
155 Thus, a wide array of data summarising the many objective measurements from each of the
156 farmlets were extracted from the Cicerone database (Scott *et al.* 2012c) from both early and late in
157 the trial period to allow direct comparisons to be made between each of the farmlets according to
158 multiple criteria. Where appropriate, the values presented over the later period have been averaged
159 over two or more years, in order to reduce the effects of year-to-year variation in some of the data
160 and thereby provide more robust measurements of the observed differences.

161 In each of the tables presented, to facilitate comparisons between farmlets, the proportion of the
162 maximum or minimum value, depending on whether a high or low result is considered desirable,

163 have been added to normalise all of the data to a scale of 0-1. The proportions were calculated as
164 described by Scott *et al.* (2000). In brief, if the desired value of a particular factor is *high* (e.g.
165 wool cut), then its average value for a particular farmlet was divided by the *maximum* of the values
166 over the three farmlets for that parameter, to give a proportion of the maximum value attained for
167 each farmlet. Alternatively, if the desired value of a parameter is *low* (e.g. number of drenches),
168 then the *minimum* value over the three farmlets was divided by the value for each farmlet, to give a
169 proportion of that minimum value for each farmlet. For example, in a case with a *high* desired
170 value, such as soil nitrogen (N), row 1 of Table 1 shows values in 2001 for farmlets A, B and C of
171 17.4, 5.2 and 13.4 respectively. Dividing each of these numbers by the maximum of the three
172 (17.4) gave proportions of 1.00, 0.30 and 0.77 respectively. In cases where one or more
173 observation was negative (e.g. cash position), the data for all three farmlets were first adjusted to
174 be equal to or above zero by adding the absolute value of the most negative observation, before
175 calculating the proportion. The overall average index values for each farmlet are simple averages
176 of all calculated indices without any attempt to weight different parameters as, in our view, any
177 weightings would be too subjective.

178 As all of the comparisons made in this paper are between average measurements from each of
179 the three unreplicated farmlets, no statistical analyses have been reported here. The case for
180 drawing causal inference between these three farming system treatments has been discussed in
181 detail by Murison and Scott (2012). Wherever feasible, the component papers of this Special Issue
182 have reported on the various statistical analyses conducted and the significance of the differences
183 found between treatments for the particular measured parameters.

184 **Results**

185 The average values of raw data derived over a wide array of parameters for each farmlet and the
186 proportions, or indices, from early (2000-2001) and late in the trial (2003-2006) are presented in
187 Table 1 and Table 2 respectively. The overall average index across all measured parameters early
188 in the trial was 0.91, 0.86 and 0.86 for farmlets A, B and C respectively (Table 1) which suggests
189 that all three farmlets were quite similar at that time. By late in the trial, the differences between

190 farmlet A and the other two farmlets had increased such that the average index across all
191 parameters was 0.91, 0.76 and 0.76 for A, B and C respectively (Table 2).

192 (insert Table 1 near here)

193 (insert Table 2 near here)

194 The relative proportions for each measured parameter are also shown graphically in Figure 1
195 and Figure 2 from both early and late in the trial. In general, the relative indices shown in Figure 1
196 (from 2000-2001) display considerable similarity among the three farmlets except for several
197 parameters which changed quickly following the imposition of differential treatments on the
198 farmlets from July 2000 (e.g. soil fertility measurements and the lower cash position of farmlets A
199 and C due to investments in pastures, fertiliser and fencing and water infrastructure).

200 (insert Figure 1 near here)

201 Figure 2 shows that the differences which developed over time between the three farmlets were
202 more marked, especially between farmlet A and the other two farmlets.

203 (insert Figure 2 near here)

204 As it is not feasible in a single paper to satisfactorily discuss all of the component issues in
205 depth, the reader is referred to those related papers noted in Tables 1 and 2 for more detailed
206 background and discussion of particular factors measured in the farmlet experiment. The most
207 significant relationships between the relative measures of the various parameters on each farmlet
208 are discussed below.

209 **Discussion and conclusions**

210 An interpretative discussion of some of the results is given below, followed by a broader, more
211 integrative discussion of the relevance of the experiment and its findings and finally, a statement of
212 the conclusions reached.

213 *Interpretation of the results*

214 *Soil nutrient levels.* Soil nutrient levels diverged quickly (Table 1) and became more different over
215 time in response to treatment (Table 2). Soil phosphorus and at times soil sulfur, were significantly
216 correlated with positive changes in botanical composition (Guppy *et al.* 2012; Shakhane *et al.*

217 2012b), including sown perennial grasses and legumes, pasture quality (Shakhane *et al.* 2012a),
218 liveweight and stocking rate (Hinch *et al.* 2012a). These results are consistent with other Northern
219 Tableland results which have shown the importance of improving livestock production through
220 higher rates of nutrient cycling and retention of those nutrients by livestock in the grazing
221 ecosystem without leakage below the pasture root zone (Chen *et al.* 2002). However, it is
222 noteworthy that there may not have been sufficient time for the nutrient flows within the farmlet
223 trial to have stabilised, as called for by van Keulen *et al.* (2000).

224 *Botanical composition.* The maintenance of sown perennial grasses, the moderate increase in
225 legume content and decrease in warm season grasses were closely associated with the pasture
226 renovation and increased soil fertility on farmlet A (Shakhane *et al.* 2012b). The proportion of
227 sown perennial grasses was maintained somewhat better by intensive rotational grazing (farmlet C)
228 than under typical management (farmlet B) which had a substantial decline in this group as well as
229 a large increase in warm season grasses.

230 Kemp (2000) described how grazing management tactics need to be used that encourage the
231 more desirable species to persist. He stated that controlling the ‘ability of animals to select what
232 they eat’ is an essential part of grazing management as continuous grazing, especially of large
233 areas, leads to patches developing in pastures. As found by Shakhane *et al.* (2012b), the
234 combination of extended grazing periods and moderate inputs resulted in a greater level of patch
235 grazing on farmlet B.

236 The importance of legumes in grazed pastures in Australia is well known. Even though there
237 was, in general, a low legume content (mostly white clover) on all the farmlets, this was associated
238 with the drier-than-average seasons experienced (Behrendt *et al.* 2012c). Nevertheless, farmlet A
239 had significantly higher legume content than either of the other farmlets due to the dual effects of
240 higher soil fertility and its longer graze and shorter rest periods than farmlet C (Shakhane *et al.*
241 2012a). Singh *et al.* (1999) have noted that higher soil phosphorus levels tend to be associated
242 with higher legume content in pastures, even during dry periods. The persistence of the perennial
243 legume, white clover, on the Northern Tablelands of NSW, was studied in a long-term trial over

244 more than 30 years by Hutchinson *et al.* (1995). When the vegetative presence of the legume was
245 reduced to low levels, they found that it was difficult to get substantial recruitment from seed
246 pools, resulting in low levels of legume especially following periods of drought (Hutchinson *et al.*
247 1995). As reported by McCaskill and Blair (1988), in dry seasons, legume growth tends to be
248 consumed rather than accumulated, resulting in increased animal liveweights. Even though the
249 levels of legume measured on the Cicerone farmlet experiment were generally low, they still had
250 significant effects on livestock production (Hinch *et al.* 2012a) and wool growth (Cottle *et al.*
251 2012).

252 *Herbage mass and quality.* In general, farmlets B and C had much higher levels of dead and total
253 herbage than farmlet A whereas all three farmlets tended to have similar levels of green herbage.
254 Although farmlet C reported similar levels of green herbage, its availability to the grazing animal
255 was less due to the intensive nature of its grazing management which meant that a much smaller
256 proportion of the farmlet was accessible to the livestock at any one time. With its greater level of
257 temperate species (Shakhane *et al.* 2012b) and higher soil fertility, farmlet A had significantly
258 higher digestibility levels resulting in a higher level of green digestible herbage over much of the
259 experiment (Shakhane *et al.* 2012a). Digestible herbage is known to be increased substantially by
260 higher levels of soil P and S resulting in increased animal production (Saul *et al.* 1999).

261 *Potential pasture growth.* Measured pasture growth was similar between farmlets (Shakhane *et al.*
262 2012a), due to the generally dry conditions. However, the level of greenness, detected by
263 measuring the normalised difference vegetation index (NDVI) via Landsat satellite images, which
264 is a surrogate measure of potential pasture growth, was significantly higher for farmlet A compared
265 to the two other farmlets (Donald *et al.* 2012). Pasture growth rate is known to be the principal
266 factor supporting changes in stocking rate which diverged over time between the farmlets (Hinch *et*
267 *al.* 2012a).

268 *Liveweight.* The substantial differences between animal performance of ewes, hoggets, wethers
269 and cattle were linked to the level of green digestible herbage and the degree of dietary choice
270 offered under the different grazing management regimes (Hinch *et al.* 2012a). Under the intensive

271 rotational grazing regime, high stocking densities meant that animals competed intensely for the
272 green digestible herbage available as shown by the rapid disappearance rate of the green
273 component of pastures on that farmlet (Shakhane *et al.* 2012c). The differences between animal
274 liveweights on farmlets were greatest when the period of grazing rest on farmlet C was longest.
275 Liveweights were also affected by the amount of legume herbage, stocking rate and the amount of
276 supplement fed (Hinch *et al.* 2012a).

277 *Fat scores and reproduction.* As with animal liveweights, the fat scores of breeding ewes tended
278 to be higher on the two farmlets that offered longer graze periods (A and B) (Hinch *et al.* 2012b).
279 The greater levels of green digestible herbage and the flexible grazing regime also led to higher
280 levels of pregnancy in scanned ewes of farmlets A and B (Hinch *et al.* 2012b).

281 *Wool production and quality.* The differences between farmlets in wool production per head and
282 wool quality whilst often significant, were not large. However, due to the combination of higher
283 production per head and stocking rate, the amount and value of wool production per hectare was
284 substantially greater on farmlet A compared to the other farmlets (Cottle *et al.* 2012).

285 *Parasites.* Overall, gastrointestinal nematodosis (GIN), as indicated by faecal worm egg count,
286 was significantly reduced on farmlet C compared with the other two farmlets (Colvin *et al.* 2008;
287 Walkden-Brown *et al.* 2012). The greatest impact was seen on *Haemonchus contortus*, with a
288 lesser impact on the other major nematode species (*Trichostrongylus* spp. and *Teladorsagia*
289 *circumcincta*). As a consequence, sheep on farmlet C received fewer anthelmintic treatments.

290 Lighter liveweights and fleece weights per head were observed on sheep from farmlet C, especially
291 early in the trial when the grazing rest periods were longest. Comparisons of GIN-free sheep
292 (treated with long-acting anthelmintics) and sheep with natural infections showed that there was no
293 impact of GIN on production traits on farmlet C whereas, bodyweight, fat score, fleece weight and
294 pregnancy rate were all higher in worm-free sheep on farmlets A and B. There was strong
295 evidence that the control of GIN on farmlet C was mediated through the intensive grazing
296 management which interrupted the free-living stages of the nematode life-cycle (Colvin *et al.*
297 2012). Short grazing periods (2-4 days) prevented autoinfection and sufficient rest periods

298 prevented large scale re-infection by allowing a significant decline of infective larval populations
299 on pasture.

300 *Economic outcomes.* The modelling which complemented the experimental results of the Cicerone
301 experiment has been discussed in other papers in this issue (Behrendt *et al.* 2012a; Behrendt *et al.*
302 2012b; Behrendt *et al.* 2012c; Scott *et al.* 2012a; Scott *et al.* 2012b). Modelling was especially
303 important in the economic analyses, given the run of poor years experienced over most of the
304 experimental period. The economic analyses used different strategies to answer a sequence of
305 related questions.

306 Scott *et al.* (2012b) used a representative-farm approach to provide realistic ‘full-scale’
307 financial results by adjusting the results from the farmlet scale to that of a commercial-scale farm
308 representative of the study area. This allowed the analysis to be extended from simple enterprise
309 gross margins to the more useful cash flow analysis for a farm as a whole. A comparison of the
310 two approaches showed the importance of understanding the meaning of the financial measures
311 used. In terms of gross margins, the full-scale Farm A performed considerably better than Farm B
312 or Farm C. Farm A also produced more wool and beef per hectare than Farm B or Farm C. But in
313 terms of cash flow, Farm B was far superior to Farm A because the latter was not able to cover its
314 high level of fixed costs, due to its high investments in pastures and soil fertility within the
315 duration of the trial and thus experienced negative cash balances for most of the period 2000-2005.
316 Farm C was inferior to Farm B in business returns due to a number of factors such as the grazing
317 rest period being too long, with high stocking densities resulting in lower liveweights for both
318 sheep and cattle and lower wool cut per head for sheep. The below-average rainfall conditions
319 experienced during 2000-2006 meant that direct economic analysis of the experimental results did
320 not include the potentially higher returns from average or above-average rainfall years that may
321 have allowed Farm A to pay its debt. This suggested that some modelling was needed to evaluate
322 the three farm systems under a wider range of conditions.

323 Scott *et al.* (2012a) introduced climatic variability into the analysis by using a stochastic
324 discounted cash flow model based on the Cicerone farmlet data. The analysis evaluated Farms A

325 and B at a commercial scale over 20 years, so that the return on investment could be evaluated. In
326 this case, over a period of 20 years, Farm A was found to be more profitable but also more ‘risky’
327 (variable), especially at the highest stocking rate explored (15 dse/ha). This indicated there was a
328 need to understand the trade-offs between risks and returns and led to further economic analyses.

329 Behrendt *et al.* (2012b) studied the risk-return trade-off in the context of pasture persistence and
330 fertiliser application under climatic uncertainty through a model calibrated using the results from
331 the Cicerone farmlet experiment. The study found that it was economically efficient to reduce both
332 fertiliser inputs and stocking rates with increasing fertiliser costs, which also reduced the
333 variability of returns (or the riskiness of the system). Although this strategy maintained similar
334 levels of total available pasture and per head livestock performance, it led to a reduction in the
335 persistence of desirable species within the sward, which could affect future returns. These results
336 revealed the importance of embedding risk in the decision process, so that decisions can be
337 adjusted as climatic variability unfolds.

338 Behrendt *et al.* (2012a) built upon the previous analysis by embedding risk into the decision
339 process to allow optimisation methods to distinguish between strategic decisions involving long
340 timeframes and tactical decisions that can be adjusted over the short-term. They argued that, in
341 evaluating the benefits of adopting alternative technologies, the way in which risk interacts with
342 management must be considered. Their study examined the conflicting goals of maximising profit
343 while achieving persistence of desirable species within a grazing system by deriving optimal
344 decision rules for the seasonal management of a paddock in the presence of climatic uncertainty.
345 Results showed the conditions under which a pasture should be renovated, given a grazing rest, or
346 have its stocking rate reduced, based on available pasture and its composition at the start of each
347 season. Typically, the stocking rate decision rule was driven by quantities of available pasture in
348 spring, whereas during summer, autumn and winter the decision was influenced by both the pasture
349 mass and the proportion of desirable species in the sward. The lowest stocking rates tended to
350 occur during winter and summer, with the highest during spring. As such, a seasonal savings and
351 consumption pattern was derived that was optimal for the given economic conditions. The re-

352 sowing of pastures was identified to be optimal only under severely degraded states of the pasture
353 resource, and was most prominent in autumn and winter. These findings provide a general
354 framework for evaluating the performance of grazing systems under climatic uncertainty whilst
355 taking pasture persistence into account.

356 *Climate constraints.* Details of the rainfall and temperature experienced over the experimental
357 period and their relationship to the long-term climatic record have been provided by Behrendt *et al.*
358 (2012c). In brief, the trial experienced drier-than-average soil moisture conditions and suffered
359 from more frequent and more severe frosts than average. As explained in that paper, the results
360 from the farmlet trial needed to be interpreted with that knowledge; this is the reason that the
361 outcomes from the modelling were important to understanding the findings in the context of longer
362 timeframes.

363 *Integrative discussion*

364 As large grazing experiments experience considerable variation over space and with changing
365 climatic conditions over time, one cannot expect such experiments to have great precision
366 (Spedding and Brockington 1976). Our experience accords with Tanaka *et al.* (2008) who stated
367 that research on farming systems which attempts to integrate multi-disciplinary strands can be
368 extremely difficult to fund, carry out, statistically analyse and publish. The integration of the
369 multiple findings from this research has been a challenge for all project participants including the
370 livestock producers who led this Project. Also, during the conduct of this farmlet experiment, there
371 were unavoidable compromises that had to be made to achieve a balance between practical
372 relevance and scientific rigour.

373 Whilst many farming system studies have been useful, some have failed to adequately mimic
374 the practices of commercially relevant farming systems (Thomson *et al.* 1995). In contrast, the
375 Cicerone farmlet experiment was designed and found to be highly relevant to producer members
376 (Edwards *et al.* 2012).

377 Results indicated that the relative performance of the three farmlets diverged considerably over
378 the duration of the experiment and over a wide array of criteria. The strategy of pasture renovation

379 with higher soil fertility employed on farmlet A resulted in a substantial improvement in overall
380 performance compared to farmlets B and C which were similar overall. This is consistent with the
381 findings of Waller *et al.* (2001) who found that greater increases in animal production came about
382 by upgrading pastures than through grazing management.

383 The integrated results of the Cicerone farmlet experiment were also, to a large extent, consistent
384 with the results of the national Sustainable Grazing Systems (SGS) experiment. Thus, as reported
385 by Sanford *et al.* (2003), there was a large effect of soil phosphorus, botanical composition, legume
386 content and stocking rate. Whereas the SGS trial used modelling to suggest that rotational grazing
387 was unlikely to markedly increase pasture production, the Cicerone farmlet experiment confirmed
388 this experimentally. Compared with flexible rotational grazing, intensive rotational grazing did not
389 lift pasture growth substantially over the duration of the trial and resulted in generally lower per
390 head and per hectare animal production. In another SGS study, Chapman *et al.* (2003) concluded
391 that 'neither grazing method explored will optimise system performance under all conditions'
392 because rigid grazing rules impact on both pastures and animals. Overall, they concluded that
393 graziers need to strive for both high per-head animal performance as well as high perennial grass
394 persistence in order to capture the growth and environmental benefits attributed to that pasture
395 component. They suggested that farms might employ more tactical grazing strategies that combine
396 set stocking and rotational systems on different parts of their farms at different times, in order to
397 ensure high per head performance.

398 The Cicerone experiment compared different grazing management systems to those studied by
399 Chapman *et al.* (2003). Whereas they compared low and high rates of fertiliser application (6.4
400 and 25 kg P/ha/year respectively), the Cicerone farmlet experiment was managed to achieve either
401 moderate or high levels of soil fertility, based on soil tests. For comparison, the annual rates of
402 fertiliser applied to farmlets B and C were approximately 4.9 and 4.3 kg P/ha respectively whereas
403 farmlet A received approximately 13.1 kg P/ha/year (Guppy *et al.* 2012). It is noteworthy that the
404 'higher' level of P application employed on farmlet A was not 'high' compared to many other
405 studies of pasture fertilisers in Australia.

406 In a study of different grazing strategies in northern Queensland, O'Reagain *et al.* (2011) found
407 that, under variable climatic conditions, the best financial and sustainable outcomes over the long-
408 term were achieved through a variable stocking strategy that allowed the adjustment of stocking
409 rate early in each dry season. Such a system is similar to the flexible grazing management
410 employed on farmlets A and B in this experiment. However, in view of the high costs incurred in
411 the initial years on farmlet A and the pasture degradation that developed over time on farmlet B, it
412 is clear that decisions need to be made that retain desirable pasture species in the pasture in such a
413 way that the life of a pasture is extended towards a region of optimal condition and management
414 (Behrendt *et al.* 2012a).

415 In a replicated grazing study of native pastures on the NW Slopes of NSW, Lodge *et al.* (2003)
416 found that, compared with continuous grazing, rotational grazing or the addition of legume and
417 fertilizer resulted in substantial improvements to the production and sustainability of those
418 pastures. Research conducted on a site adjacent to the Cicerone farmlet experiment found that a
419 combination of a persistent legume (white clover) and a strongly perennial temperate grass
420 (phalaris) had considerable sustainability benefits. Thus, McLeod *et al.* (2006) found that such a
421 pasture had a deeper rooting depth which enabled more soil water extraction which was associated
422 with greater sustainability characteristics (Scott *et al.* 2000) than either a phalaris-dominant pasture
423 or a degraded pasture. Also, on the Central Tablelands of NSW, sown perennial grass pastures
424 were found to contribute to sustainability, provided that they last long enough to provide positive
425 economic returns (Dowling *et al.* 2006).

426 We acknowledge that, due to a lack of resources (Scott *et al.* 2012c), insufficient measurements
427 relating to sustainability, such as the indirect effects of the state of pastures on ground-water, soil
428 acidification and erosion over the long-term, were undertaken within the farmlet experiment. As
429 noted by Jones *et al.* (1995) and Saul and Chapman (2002), there remains a need to evaluate the
430 environmental effects of grazed systems over the long-term as it can take many years for pasture
431 composition and soil fertility to reach steady states.

432 When seeking answers to broad agroecological questions such as agricultural sustainability,
433 Edwards *et al.* (1993) pointed out the need for multi-disciplinary approaches which not only
434 include farmers but also take a ‘whole farm level’ approach. Modelling of livestock farming
435 systems has also been suggested as useful in striving for more sustainable systems (Gibon *et al.*
436 1999) so long as it is conducted in an inter-disciplinary fashion. The Cicerone farmlet experiment
437 adopted both multi-disciplinary and modelling approaches.

438 Grazed experiments which have comprehensively measured most components of a farming
439 system are rare. Saul *et al.* (2011) found that upgraded pastures were associated with an increased
440 stocking rate, higher ewe liveweights, condition scores, lambing and weaning percentages, wool
441 cut per head and fibre diameter but with no increase in supplementary feeding. They also
442 estimated that the break-even point for financial returns on investments in upgraded pastures
443 occurred after 7 years. In addition, they estimated an internal rate of return on pasture
444 improvement of 27%, assuming the upgraded pastures were managed to persist for 12 years. In the
445 case of pasture renovation on the Cicerone farmlet experiment, Behrendt *et al.* (2006) estimated
446 that the risk efficient rate of pasture replacement was 4% per annum.

447 Over the duration of the trial, the level of green herbage declined well below critical thresholds
448 on a number of occasions, as shown by pasture assessments (Shakhane *et al.* 2012a) and the trends
449 in liveweights and supplement fed (Hinch *et al.* 2012a). The ability of sheep to select a high
450 quality diet decreases as the level of green herbage on offer declines below the critical threshold of
451 550 kg green DM/ha (Hamilton *et al.* 1973). Levels of green herbage were affected by rainfall and
452 season, pasture species, soil fertility, stocking rate and grazing management. Thus, the low levels
453 of green herbage on farmlet A (Shakhane *et al.* 2012a) were exacerbated by the stocking rate being
454 too high at times, as evidenced by the high levels of supplement required in the later years of the
455 trial. Had a greater level of green herbage been retained, it is likely that pasture growth and
456 persistence would have been enhanced with less supplement required (Shakhane *et al.* 2012a).

457 Both farmlets B and C recorded higher levels of dead herbage than farmlet A (Shakhane *et al.*
458 2012a). On farmlet B, the low levels of green herbage in winter were related to the rise of warm

459 season grasses and low levels of legume. Farmlet C also experienced similar low levels of green
460 herbage and digestibility to those on farmlet B but, because only a small proportion of the farmlet
461 was available to the grazing animals at any one time, the high stock density resulted in intense
462 competition for and rapid depletion of green herbage during each brief grazing event (Shakhane *et*
463 *al.* 2012c). Chapman *et al.* (2003) have argued that the rigid application of grazing management
464 regimes can result in pastures of lower quality. In rotationally grazed systems, there are
465 compromises and trade-offs which occur between pasture and animal production such as the lower
466 legume levels and lower levels of per head and per hectare production (Saul and Chapman 2002).

467 Hall *et al.* (2011) compared three grazing systems, from continuous grazing through to
468 intensive grazing, under extensive commercial conditions in Queensland and found that diet
469 quality tended to be lower in the more intensive grazing systems. Others too have found that
470 longer grazing rest periods can result in lower quality herbage because of a higher proportion of
471 stem to digestible leaf (Waller *et al.* 2001).

472 *Critique of the three farmlet management systems*

473 The control farmlet (B), which was managed according to guidelines considered typical by the
474 region's livestock producers, was found to have the highest cash position at the end of the trial.
475 This was achieved largely by accumulating income through employing a modest stocking rate and
476 constraining expenditure. In doing so, it achieved high per head weight gains and performed best
477 in terms of lambs marked per ewe joined and in lamb mortality. However, in other respects it was
478 found wanting, especially in its pastures, which became degraded over time, with low levels of
479 legume, declining levels of sown perennial grasses, increasing levels of warm season grasses and
480 evidence of patch grazing and increased broadleaf weeds in all paddocks except one which was
481 renovated in 2004 (Shakhane *et al.* 2012b). Thus, in spite of its superior financial position, we
482 contend that this farmlet represents the least sustainable management option.

483 In contrast, farmlet A, which benefited substantially from renovated pastures and higher soil
484 fertility, suffered financially from the high rate of pasture renovation (29% of the farmlet in the
485 first year). This was largely an artefact of Project management which determined that the farmlets

486 had to be developed quickly so that differences between farmlets would emerge over a short
487 timeframe. Farmlet A also suffered financially due to the cost of sowing two paddocks of so-
488 called ‘high performance’ pastures based on Italian ryegrass, which producer members wanted
489 investigated. In both cases, the Italian ryegrass pastures failed to persist longer than 18 months and
490 needed to be re-sown to perennial pastures.

491 As noted by Carter and Day (1970), there needs to be a sufficient financial incentive if
492 producers are to consider a more productive strategy, such as increased stocking rates, with its
493 inherent risks and need for greater managerial skill. The modelling (Behrendt *et al.* 2006) and
494 economic risk (Scott *et al.* 2012a) studies conducted, confirmed that optimal solutions demand that
495 pastures be managed to persist over long-periods in order to justify investment in such
496 technologies. Thus, assuming a rate of pasture renovation at the most risk efficient level of 4% per
497 annum (Behrendt *et al.* 2006) and a stocking rate of either 11.9 or 15 dse/ha, farmlet A was shown
498 to have the potential for a substantially higher cumulative net present value over a 20 year horizon
499 than an analysis based on the artificially high rate of pasture renovation during this trial (Scott *et*
500 *al.* 2012a); the risk level for the 11.9 dse/ha scenario was substantially lower than for the 15 dse/ha
501 scenario. As pointed out by Scott *et al.* (2012a), given some better seasons, this farmlet had much
502 more potential for ‘upside risk’, or more favourable economic outcomes, compared to either of the
503 other farmlets.

504 In terms of meeting some criteria which are associated with ‘sustainability’ (Scott *et al.* 2000),
505 farmlet A achieved higher levels of soil nutrients, temperate perennial grasses, legumes and higher
506 per hectare livestock production than the other farmlets whilst also achieving high levels of animal
507 production on a per head basis. However, it must be acknowledged that these increases came at a
508 considerable cost and that insufficient environmental factors were able to be measured. Thus,
509 although farmlet A reached a substantially higher average index (0.91) over all measured
510 parameters by the latter half of the experiment compared to farmlets B and C (0.76 and 0.76
511 respectively), it is difficult to conclude that any one system would be likely to be more profitable
512 or sustainable over the long-term.

513 Regarding the higher numbers of gastrointestinal nematodes on both farmlets A and B
514 compared to farmlet C (Walkden-Brown *et al.* 2012), it is clear that the two farmlets which
515 employed flexible rotational grazing were constrained substantially by needing to graze multiple
516 mobs across no more than 8 paddocks per farmlet. No doubt, better control could have been
517 achieved if more paddocks had been allowed and if more deliberate use had been made of cattle to
518 graze paddocks before sheep, as recommended by Niezen *et al.* (1996). The question therefore
519 arises, how many paddocks would be optimal for such flexible rotational grazing systems?
520 Through modelling, Morley (1968) predicted that, from the point of view of pasture growth alone,
521 the optimum number of paddocks for a rotational system would probably be less than 10. In view
522 of the experience of this farmlet trial, with its desire of taking into account a wider array of factors,
523 including multiple mobs and control of internal parasites, we suggest that it is likely that the
524 optimum number of paddocks would be well above 10 per farm.

525 Whilst the intensive rotational grazing regime of farmlet C clearly resulted in superior control
526 of gastrointestinal nematodes, this management regime did not result in increased pasture
527 production, higher soil phosphorus, higher stocking rates or higher profit. Whilst the differences
528 between farmlets in green herbage were not significant, the digestibility of both green and dead
529 herbage was significantly lower on farmlet C (and B) than on farmlet A. Nevertheless, farmlet C
530 had a slightly finer wool fibre diameter, slightly stronger staple strength for ewes, needed less
531 supplementary feeding and had slightly higher ground cover. Due to the nature of intensive
532 grazing systems, a much smaller proportion of paddocks on farmlet C was grazed at any one time;
533 this higher stock density during grazing is likely to be the main reason for the lower animal
534 performance on farmlet C as the animals had to compete intensely for the available green herbage.
535 It is also likely that part of the reason for the lower animal performance on farmlet C was the low
536 levels of legume on this farmlet which were associated with low soil nitrogen levels which would
537 in turn have constrained pasture growth in winter.

538 It is interesting to speculate how the performance of farmlet C might have been enhanced. One
539 suggestion from a Cicerone member, late in the trial period, was that a 'higher input' version of

540 farmlet C (similar to the inputs on farmlet A) should be studied (Edwards *et al.* 2012).
541 Unfortunately, no funding was available to create this fourth farmlet to allow such a comparison.
542 As with the other farmlets, it would also have been interesting to observe and measure the
543 consequences of some better seasons on farmlet C though we contend that the moderate soil
544 fertility, the low level of legumes and the moderate proportion of temperate species on this farmlet
545 would have limited the potential for high pasture growth rates.

546 *Implications for optimal management of grazing enterprises*

547 In spite of the trial's limited duration of 6.5 years and the constraints imposed by the drier-than-
548 average conditions, the consideration of the multiple lines of accumulated evidence, together with
549 modelling and optimisation procedures, has yielded a number of important outcomes from this
550 body of work. Whereas modelling analyses suggested that farmlet A had the greater *potential* for
551 profit over the long-term, given a more representative climatic experience, one needs to reflect on
552 how that might be achieved with lower levels of risk. If, for the creation of optimal net worth
553 outcomes, pastures need to be maintained to persist over some 25 years (equivalent to a 4% per
554 annum replacement rate) this would mean that managers would need to pay much greater attention
555 to the maintenance of soil fertility and the strategic resting of pastures before they reach a critical
556 degraded state (Behrendt *et al.* 2012a).

557 The challenge for graziers is to 'simultaneously balance many balls in the air' (Williams 1994).
558 Based on the interacting factors described in this paper, we suggest it is important to distinguish
559 between the short-term, tactical 'balls' and the longer-term strategic 'balls'. Some tactical
560 decisions include moving stock between paddocks, supplementary feeding and drenching. In
561 contrast, strategic decisions can be crucial for delivering optimal results over the long-term. Such
562 decisions include: maintaining soil fertility and an adequate level of legumes through nutrition and
563 grazing management to help manipulate pasture composition, the gradual renovation of pastures to
564 enhance the proportion of desirable species, integrated parasite management, allowing increases in
565 stocking rate only when the potential for pasture growth has been enhanced, the creation of more

566 protection for lambing ewes, and the provision of sufficient paddocks to facilitate rotation when
567 necessary.

568 Decisions regarding changes to stocking rate would be aided greatly by being able to regularly
569 monitor all paddocks through remote sensing in order to estimate potential pasture growth rate
570 (Donald *et al.* 2012) which is a fundamental parameter governing stocking rates. Regular and
571 timely estimates of paddock and farm-scale green herbage mass, which can also be derived from
572 satellite images (Edirisinghe *et al.* 2011), would also greatly assist graziers to ensure that critical
573 levels of green herbage are always available to their stock as a means of optimising pasture
574 utilisation whilst avoiding over-grazing which can threaten pasture persistence and limit the need
575 for supplementary feeding.

576 In addition, regular monitoring of liveweights and condition scores would enable the detection
577 of livestock performance and allow changes of stocking rates or paddock moves to be made in a
578 more timely way. Options for reducing worm burdens in sheep within flexible rotational grazing
579 systems include preparation of ‘clean’ pastures for young, susceptible, stock (Bailey *et al.* 2009),
580 mixed grazing with cattle or sheep of lesser susceptibility (wethers), and lightly grazing pasture
581 with cattle prior to grazing with sheep.

582 Ultimately, improved tools need to be developed to assist graziers to work their way through
583 these decision dilemmas. There is a need for more robust and timely delivery of optimal solutions
584 which take into account the different rates of change of the various factors which can be controlled
585 by management so that long-term profits can be realised without damage to the natural resources
586 which support grazing enterprises.

587 This paper has highlighted the complexity of managing grazing enterprises and the problems of
588 making any simple, prescriptive recommendations. Grazed farms are challenging agroecosystems
589 that require a wide range of measurements, skills and observations that need to be continually
590 reviewed to achieve desired outcomes. In addition, conditions and circumstances vary from
591 paddock to paddock, from farm to farm, whilst skill levels and goals vary from one livestock
592 producer to another. Issues that might appear to be simple – such as maintaining a desirable

593 pasture composition – are affected not only by the species and their relative growth rates across
594 different seasons but also by soil type, soil fertility, slope, aspect, rainfall and temperature, grazing
595 rest, livestock species, livestock class, ground cover and the manager’s attitude to risk.

596 The particular version of intensive rotational grazing tested here (on farmlet C) provided
597 substantial benefits in terms of animal health but the level of animal production supported by this
598 system was overly constrained by the limited dietary choice offered grazing animals and the low
599 levels of legume which restricted the protein supply for livestock. The typically managed farmlet
600 (B), which had the best cash position over the relatively short duration of this trial, was found to
601 have developed several negative attributes which suggests that it may not be a profitable or
602 sustainable alternative management system over the long-term. In answer to the hypothesis put
603 earlier in this paper, we conclude that more profitable and sustainable outcomes are most likely to
604 arise from grazing enterprises which are proactively managed towards optimal outcomes which
605 include the maintenance of sufficient desirable perennial grasses, combined with adequate legume
606 content and are supported by the maintenance of soil fertility whilst employing flexible rotational
607 grazing.

608 **Acknowledgements**

609 This Project was generously funded by Australian Wool Innovation with additional financial
610 support from the Australian Sheep CRC and the University of New England. It benefited
611 substantially from the substantial in-kind contributions of NSW Department of Primary Industries,
612 CSIRO Livestock Industries, Betty Hall Pty Ltd and the University of New England. Whilst too
613 numerous to mention individually here, the livestock producer members of the Cicerone Project
614 were generous in their support. We are indebted to those many collaborators who produced the
615 results which we have summarised from the component papers of this Special Issue.

616 **References**

617 Andrew MH, Lodge GM, Mason WK, Price RJ (2003) The Sustainable Grazing Systems National
618 Experiment. 2. Scientific outcomes and effectiveness of the research and development processes.
619 *Australian Journal of Experimental Agriculture* **43**, 993-1013.

620 Bailey JN, Walkden-Brown SW, Kahn LP (2009) Comparison of strategies to provide lambing
621 paddocks of low gastro-intestinal nematode infectivity in a summer rainfall region of Australia.
622 *Veterinary Parasitology* **161**, 218-231.

623 Behrendt K, Cacho O, Scott JM, Jones R (2006) Methodology for assessing optimal rates of
624 pasture improvement in the high rainfall temperate pasture zone. *Australian Journal of*
625 *Experimental Agriculture* **46**, 845-849.

626 Behrendt K, Cacho O, Scott JM, Jones R (2012a) Optimising pasture and grazing management
627 decisions on the Cicerone Project farmlets over variable time horizons. *Animal Production Science*
628 **52**, (in press).

629 Behrendt K, Scott JM, Cacho O, Jones R (2012b) Simulating the impact of fertiliser strategies and
630 prices on the economics of developing and managing the Cicerone Project farmlets under climatic
631 uncertainty. *Animal Production Science* **52**, (in press).

632 Behrendt K, Scott JM, Mackay D (2012c) Assessing the climate experienced during the Cicerone
633 farmlet experiment against the climatic record. *Animal Production Science* **52**, (in press).

634 Briske DD, Sayre NF, Huntsinger L, Fernandez-Gimenez M, Budd B, Derner JD (2011) Origin,
635 Persistence, and Resolution of the Rotational Grazing Debate: Integrating Human Dimensions Into
636 Rangeland Research. *Rangeland Ecology & Management* **64**, 325-334.

637 Carter E, Day H (1970) Interrelationships of stocking rate and superphosphate rate on pasture as
638 determinants of animal production. I. Continuously grazed old pasture land. *Australian Journal of*
639 *Agricultural Research* **21**, 473-491.

640 Cawood M (2004) Working with Nature on Lana In 'The Land'. July 26, 2004.

641 Cawood M (2011) Grazing systems study "missed the point". In 'The Land'. December 1, 2011.

642 Chapman DF, McCaskill MR, Quigley PE, Thompson AN, Graham JF, Borg D, Lamb J, Kearney
643 G, Saul GR, Clark SG (2003) Effects of grazing method and fertiliser inputs on the productivity
644 and sustainability of phalaris-based pastures in Western Victoria. *Australian Journal of*
645 *Experimental Agriculture* **43**, 785-798.

646 Chen W, Scott J, Blair G, Lefroy R, Hutchinson K, King K, Harris C (2002) Diet selection and
647 productivity of sheep grazing contrasting pastures. *Australian Journal of Agricultural Research* **53**,
648 529-539.

649 Colvin AF, Walkden-Brown SW, Knox MR (2012) Role of host and environment in mediating
650 reduced gastrointestinal nematode infections in sheep due to intensive rotational grazing.
651 *Veterinary Parasitology* **184**, 180-192.

652 Colvin AF, Walkden-Brown SW, Knox MR, Scott JM (2008) Intensive rotational grazing assists
653 control of gastrointestinal nematodosis of sheep in a cool temperate environment with summer-
654 dominant rainfall. *Veterinary Parasitology* **153**, 108-120.

655 Cook SJ, Lazenby A, Blair GJ (1976) Comparative responses of *Lolium perenne* and *Bothriochloa*
656 *macra* to temperature, moisture, fertility and defoliation. *Australian Journal of Agricultural*
657 *Research* **27**, 769-778.

658 Cook SJ, Lazenby A, Blair GJ (1978) Pasture degeneration. 1. Effect on total and seasonal pasture
659 production. *Australian Journal of Agricultural Research* **29**, 9-18.

660 Cottle D, Gaden CA, Hoad J, Scott JM, Lance D (2012) Evaluation of the impact of pasture and
661 grazing management on wool production, fibre diameter profiles, wool quality and wool returns.
662 *Animal Production Science* **52**, (in press).

663 Donald GE, Scott JM, Vickery PJ (2012) Satellite derived evidence of whole farmlet and paddock
664 responses to management and climate. *Animal Production Science* **52**, (in press).

665 Dowling PM, Kemp DR, Ball PD, Langford CM, Michalk DL, Millar GD, Simpson PC,
666 Thompson RP (2005) Effect of continuous and time-control grazing on grassland components in
667 south-eastern Australia. *Australian Journal of Experimental Agriculture* **45**, 369-382.

668 Dowling PM, Michalk DL, Kemp DR, Millar GD, Priest SM, King WM, Packer IJ, Holst PJ,
669 Tarleton JA (2006) Sustainable grazing systems for the Central Tablelands of New South Wales. 2.
670 Effect of pasture type and grazing management on pasture productivity and composition.
671 *Australian Journal of Experimental Agriculture* **46**, 457-469.

672 Earl J, Jones C (1996) The need for a new approach to grazing management - Is Cell grazing the
673 answer? *The Rangeland Journal* **18**, 327-350.

674 Eberhardt LL, Thomas JM (1991) Designing environmental field studies. *Ecological Monographs*
675 **61**, 53-73.

676 Edirisinghe A, Hill MJ, Donald GE, Hyder M (2011) Quantitative mapping of pasture biomass
677 using satellite imagery. *International Journal of Remote Sensing* **32**, 2699-2724.

678 Edwards C, Gaden CA, Marchant R, Coventry T, Dutton P, Scott JM (2012) Delivering extension
679 and adult learning outcomes from the Cicerone Project by 'comparing, measuring, learning and
680 adopting'. *Animal Production Science* **52**, (in press).

681 Edwards CA, Grove TL, Harwood RR, Colfer CJP (1993) The Role of Agroecology and Integrated
682 Farming Systems in Agricultural Sustainability. *Agriculture Ecosystems & Environment* **46**, 99-
683 121.

684 Gibon A, Sibbald AR, Flamant JC, Lhoste P, Revilla R, Rubino R, Sorensen JT (1999) Livestock
685 fanning systems research in Europe and its potential contribution for managing towards
686 sustainability in livestock farming. *Livestock Production Science* **61**, 121-137.

687 Guppy CN, Scott JM, Edwards E, Blair GJ (2012) Whole-farm management of soil nutrients drives
688 productive grazing systems: the Cicerone farmlet experiment confirms earlier research. *Animal*
689 *Production Science* **52**, (in press).

690 Hacker R (1993) A brief evaluation of time control grazing. In 'Proceedings of the Eighth Annual
691 Conference of the Grasslands Society of NSW'. Orange, NSW pp. 82-89.

692 Hall TJ, McIvor J, Jones P, MacLeod N, McDonald C, Reid D, Smith D, Delaney K (2011) Final
693 report: Volume I - Investigating Intensive Grazing Systems in Northern Australia. (Meat &
694 Livestock Australia Limited).

695 <http://www.mla.com.au/CustomControls/PaymentGateway/ViewFile.aspx?GYuXLNalmke0hkUG>
696 [MDR0Hq7ke55w29oGRr/R/qg3AoJsOM005VLFzKnPrSxthKQT3EYMKKAfsht7d1Tnt3BqiA==](http://www.mla.com.au/CustomControls/PaymentGateway/ViewFile.aspx?MDR0Hq7ke55w29oGRr/R/qg3AoJsOM005VLFzKnPrSxthKQT3EYMKKAfsht7d1Tnt3BqiA==)
697 . 96 pp.

698 Hamilton B, Hutchinson K, Annis P, Donnelly J (1973) Relationships between the diet selected by
699 grazing sheep and the herbage on offer. *Australian Journal of Agricultural Research* **24**, 271-277.

700 Hinch GN, Hoad J, Lollback M, Marchant RS, Colvin AF, Scott JM, Mackay D (2012a) Livestock
701 weights in response to three whole-farmlet management systems. *Animal Production Science* **52**,
702 (in press).

703 Hinch GN, Lollback M, Hatcher S, Hoad J, Marchant RS (2012b) Comparison of sheep fat scores
704 and reproduction as affected by three whole-farmlet management systems. *Animal Production*
705 *Science* **52**, (in press).

706 Hutchinson K, King K, Wilkinson D (1995) Effects of rainfall, moisture stress, and stocking rate
707 on the persistence of white clover over 30 years. *Australian Journal of Experimental Agriculture*
708 **35**, 1039-1047.

709 Jones RM, Jones RJ, McDonald CK (1995) Some advantages of long-term grazing trials, with
710 particular reference to changes in botanical composition. *Australian Journal of Experimental*
711 *Agriculture* **35**, 1029-1038.

712 Kemp DR (2000) Managing grassland composition with grazing. In 'Proceedings of the Bushcare
713 Grassy Landscapes Conference: Balancing Conservation and Production in Grassy Landscapes'.
714 Clare, SA. (Eds T Barlow, R Thorburn) pp. 145-152. (Environment Australia, Canberra).

715 <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.94.1734&rep=rep1&type=pdf#page=15>
716 [8](#).

717 Lazenby A, Lovett J (1975) Growth of pasture species on the Northern Tablelands of New South
718 Wales. *Australian Journal of Agricultural Research* **26**, 269-280.

719 Lodge G (2002) Can graziers assess sustainable and productive pastures? In 'Proceedings of the
720 17th Annual Conference of the Grasslands Society of NSW'. Forbes. (Ed. AM Bowman) pp. 63-64.
721 (The Grasslands Society of NSW Inc).

722 Lodge GM, Murphy SR, Harden S (2003) Effects of grazing and management on herbage mass,
723 persistence, animal production and soil water content of native pastures. 1. A redgrass-wallaby

724 grass pasture, Barraba, North West Slopes, New South Wales. *Australian Journal of Experimental*
725 *Agriculture* **43**, 875-890.

726 Lodge GM, Scott JM, King KL, Hutchinson KJ (1998) A review of sustainable pasture production
727 issues in temperate native and improved pastures. In 'Animal Production in Australia'. Armidale,
728 NSW **22**. pp. 79-89. <http://www.asap.asn.au/livestocklibrary/1998/Lodge98.PDF>.

729 McCaskill MR, Blair GJ (1988) Medium-term climatic variation of the Northern Tablelands of
730 NSW. *Mathematics and Computers in Simulation* **30**, 159-164.

731 McCosker T (2000) Cell Grazing - the first 10 years in Australia. *Tropical Grasslands* **34**, 207-
732 218.

733 McLeod MK, MacLeod DA, Daniel H (2006) The effect of degradation of phalaris plus white
734 clover pasture on soil water regimes of a Brown Chromosol on the Northern Tablelands of NSW,
735 Australia. *Agricultural Water Management* **82**, 318-342.

736 Moore RM, Barrie N, Kipps EH (1946) Continuous and rotational grazing by Merino sheep. 1. A
737 study of the production of a sown pasture in the Australian Capital Territory under three systems of
738 grazing management. CSIR, Melbourne. 7-83 pp.

739 Morley F (1968) Pasture growth curves and grazing management. *Australian Journal of*
740 *Experimental Agriculture* **8**, 40-45.

741 Murison RD, Scott JM (2012) Statistical methodologies for making causal inference from an
742 unreplicated experiment which compared three whole-farmlet management systems. *Animal*
743 *Production Science* **52**, (in press).

744 Niezen JH, Charleston WAG, Hodgson J, Mackay AD, Leathwick DM (1996) Controlling internal
745 parasites in grazing ruminants without recourse to anthelmintics - approaches, experiences and
746 prospects. *International Journal for Parasitology* **26**, 983-992.

747 Norton BE (1998) The application of grazing management to increase sustainable livestock
748 production. In 'Animal Production in Australia'. Armidale, NSW **22**. pp. 15-26.
749 <http://www.asap.asn.au/livestocklibrary/1998/MCCMONT.PDF>.

750 O'Reagain P, Bushell J, Holmes B (2011) Managing for rainfall variability: long-term profitability
751 of different grazing strategies in a northern Australian tropical savanna. *Animal Production Science*
752 **51**, 210-224.

753 Robinson G, Lazenby A (1976) Effect of superphosphate, white clover and stocking rate on the
754 productivity of natural pastures, Northern Tablelands, New South Wales. *Australian Journal of*
755 *Experimental Agriculture* **16**, 209-217.

756 Sanford P, Cullen BR, Dowling PM, Chapman DF, Garden DL, Lodge GM, Andrew MH, Quigley
757 PE, Murphy SR, King WM, Johnston WH, Kemp DR (2003) SGS Pasture Theme: effect of
758 climate, soil factors and management on pasture production and stability across the high rainfall
759 zone of southern Australia. *Australian Journal of Experimental Agriculture* **43**, 945-959.

760 Sanjari G, Ghadiri H, Ciesiolka CAA, Yu B (2008) Comparing the effects of continuous and time-
761 controlled grazing systems on soil characteristics in Southeast Queensland. *Australian Journal of*
762 *Soil Research* **46**, 348-358.

763 Saul G, Kearney G, Borg D (2011) Pasture systems to improve productivity of sheep in south-
764 western Victoria 2. Animal production from ewes and lambs. *Animal Production Science* **51**, 982-
765 989.

766 Saul GR, Chapman DF (2002) Grazing methods, productivity and sustainability for sheep and beef
767 pastures in temperate Australia. *Wool Technology and Sheep Breeding* **50**, 449-464.

768 Saul GR, Kearney GA, Flinn PC, Lescun CL (1999) Effects of superphosphate fertiliser and
769 stocking rate on the nutritive value of perennial ryegrass and subterranean clover herbage.
770 *Australian Journal of Agricultural Research* **50**, 537-545.

771 Scott F, Cacho O, Scott JM (2012a) Economic risk analysis of livestock management system
772 options. *Animal Production Science* **52**, (in press).

773 Scott F, Scott JM, Cacho O (2012b) Whole farm returns show true profitability of three different
774 livestock management systems. *Animal Production Science* **52**, (in press).

775 Scott JM, Gaden CA, Edwards C, Paull DR, Marchant RS, Hoad J, Sutherland H, Coventry T,
776 Dutton P (2012c) Selection of experimental treatments, methods used and evolution of

777 management guidelines for comparing and measuring three grazed farmlet systems. *Animal*
778 *Production Science* **52**, (in press).

779 Scott JM, Hutchinson KJ, King K, Chen W, McLeod M, Blair GJ, White A, Wilkinson D, Lefroy
780 RDB, Cresswell H, Daniel H, Harris C, MacLeod DA, Blair N, Chamberlain G (2000) Quantifying
781 the sustainability of grazed pastures on the Northern Tablelands of New South Wales. *Australian*
782 *Journal of Experimental Agriculture* **40**, 257-265.

783 Shakhane LM, Mulcahy C, Scott JM, Hinch GN, Mackay D (2012a) Pasture herbage mass, quality
784 and growth in response to three whole-farmlet management systems. *Animal Production Science*
785 **52**, (in press).

786 Shakhane LM, Murison RD, Mulcahy C, Scott JM, Hinch GN, Mackay D, Morrow A (2012b)
787 Changes in botanical composition on three farmlets subjected to different pasture and grazing
788 management strategies. *Animal Production Science* **52**, (in press).

789 Shakhane LM, Scott JM, Lord CJ, Hinch GN (2012c) Matching the supply of pasture with the
790 demand of grazing livestock. *Animal Production Science* **52**, (in press).

791 Singh DK, Sale PWG, Gourley CJP, Hasthorpe C (1999) High phosphorus supply increases
792 persistence and growth of white clover in grazed dairy pastures during dry summer conditions.
793 *Australian Journal of Experimental Agriculture* **39**, 579-585.

794 Spedding CRW, Brockington NR (1976) Guest editorial: The study of ecosystems. *Agro-*
795 *Ecosystems* **2**, 165-172.

796 Sutherland H, Scott JM, Gray D, Woolaston R (2012) Creating the Cicerone Project: seeking
797 closer engagement between livestock producers, research and extension. *Animal Production*
798 *Science* **52**, (in press).

799 Tanaka DL, Karn JF, Scholljegerdes EJ (2008) Integrated crop/livestock systems research:
800 Practical research considerations. *Renewable Agriculture and Food Systems* **23**, 80-86.

801 Thomson EF, Bahhady FA, Nordblom TL, Harris HC (1995) A Model-Farm Approach to Research
802 on Crop Livestock Integration .3. Benefits of Crop Livestock Integration and a Critique of the
803 Approach. *Agricultural Systems* **49**, 31-44.

804 van Keulen H, Aarts HFM, Habekotte B, van der Meer HG, Spiertz JHJ (2000) Soil-plant-animal
805 relations in nutrient cycling: the case of dairy farming system 'De Marke'. *European Journal of*
806 *Agronomy* **13**, 245-261.

807 Vere DT, Campbell MH (2004) Potential economic benefits to the Australian wool industry from
808 improving pastures in the hill country of eastern New South Wales. *The Rangeland Journal* **26**,
809 161-177.

810 Walkden-Brown S, Colvin A, Hall E, Knox M, Scott JM (2012) Experiences with managing sheep
811 worm infections on three different whole-farmlet management systems. *Animal Production*
812 *Science* **52**, (in press).

813 Waller RA, Sale PWG, Saul GR, Kearney GA (2001) Tactical versus continuous stocking in
814 perennial ryegrass-subterranean clover pastures grazed by sheep in south-western Victoria - 3.
815 Herbage nutritive characteristics and animal production. *Australian Journal of Experimental*
816 *Agriculture* **41**, 1121-1131.

817 Waugh W (1997) Grazing systems in practice. In 'Proceedings of the twelfth Annual Conference of
818 the Grasslands Society of NSW'. Dubbo, NSW. (Eds AM Bowman, DL Michalk) pp. 91-94. (The
819 Grassland Society of NSW Inc).

820 Whalley R, Robinson G, Taylor J (1976) General effects of management and grazing by domestic
821 livestock on the rangelands of the Northern Tablelands of New South Wales. *The Rangeland*
822 *Journal* **1**, 174-190.

823 Williams G (1994) Landcare in agriculture - a personal view. In 'Is Agriculture on a Downward
824 Spiral? Can Landcare Make a Difference? Proceedings of the Northern Tablelands Landcare
825 Seminar'. Armidale. (Ed. JC Prior) pp. 22-24. (CaLM).

826 Wolfe E, Lazenby A (1973) Grass-white clover relationships during pasture development. 2. Effect
827 of nitrogen fertilizer with superphosphate. *Australian Journal of Experimental Agriculture* **13**, 575-
828 580.

829

830

Table 1. Average data from farmlets A, B and C early in the trial (2000 to 2001) for a wide array of soil, pasture, ground cover, livestock production, animal health, farmlet productivity and financial measurements and proportions of the highest data value of each measurement, normalised on a 0-1 scale.

| Factor | Units | Aim (High/Low) | Year(s) | Average data | | | Proportion of maximum value (for aim of H)/ minimum (for aim of L) | | | Source of data from related publication |
|----------------------------------|--------------|-------------------|-----------|--------------|-------|-------|--|------|------|--|
| | | | | A | B | C | A | B | C | |
| <i>Soil layer</i> | | | | | | | | | | |
| Soil-N | mg N/kg soil | H | 2001 | 17.4 | 5.2 | 13.4 | 1.00 | 0.30 | 0.77 | (Guppy <i>et al.</i> 2012) |
| Soil-P | mg P/kg soil | H | 2001 | 29.5 | 14.5 | 21.0 | 1.00 | 0.49 | 0.71 | (Guppy <i>et al.</i> 2012) |
| Soil-S | mg S/kg soil | H | 2001 | 14.5 | 6.5 | 8.0 | 1.00 | 0.45 | 0.55 | (Guppy <i>et al.</i> 2012) |
| <i>Pasture layer</i> | | | | | | | | | | |
| Sown perennial grasses | % | H | 2000 | 51.7 | 30.5 | 43.9 | 1.00 | 0.59 | 0.85 | (Shakhane <i>et al.</i> 2012b) |
| Legume | % | H | 2000 | 1.4 | 3.7 | 3.1 | 0.38 | 1.00 | 0.86 | (Shakhane <i>et al.</i> 2012b) |
| Warm season grasses | % | L | 2000 | 14.9 | 25.6 | 21.3 | 1.00 | 0.58 | 0.70 | (Shakhane <i>et al.</i> 2012b) |
| Total herbage mass | kg DM/ha | H | 2000-2001 | 3402 | 2791 | 3350 | 1.00 | 0.82 | 0.98 | (Shakhane <i>et al.</i> 2012a) |
| Greenness (satellite NDVI) | NDVI | H | 2000 | 321 | 341 | 341 | 0.94 | 1.00 | 1.00 | (Donald <i>et al.</i> 2012) |
| <i>Livestock production</i> | | | | | | | | | | |
| Greasy fleece weight/head (ewes) | kg/hd | H | 2001 | 3.2 | 3.2 | 3.1 | 1.00 | 0.99 | 0.96 | (Cottle <i>et al.</i> 2012) |
| Fibre diameter (ewes) | microns | L | 2001 | 19.4 | 19.1 | 19.2 | 0.99 | 1.00 | 0.99 | (Cottle <i>et al.</i> 2012) |
| Staple length (ewes) | mm | H | 2001 | 60.7 | 59.5 | 60.6 | 1.00 | 0.98 | 1.00 | (Cottle <i>et al.</i> 2012) |
| Liveweight/head (ewes) | kg/hd | H | 2000-2001 | 43.1 | 43.6 | 43.3 | 0.99 | 1.00 | 0.99 | (Hinch <i>et al.</i> 2012a) |
| Fat score/head (ewes) | score (1-5) | H | 2000-2001 | 2.6 | 2.7 | 2.7 | 0.97 | 0.99 | 1.00 | (Hinch <i>et al.</i> 2012b) |
| Liveweight/head (weaners) | kg/hd | H | 2000 drop | 20.0 | 18.8 | 18.5 | 1.00 | 0.94 | 0.93 | (Hinch <i>et al.</i> 2012a) |
| Parasites (faecal egg count) | number/gram | L | 2000-2001 | 164.3 | 203.2 | 224.7 | 1.00 | 0.81 | 0.73 | (Walkden-Brown <i>et al.</i> 2012) |
| Number of drenches/mob/year | number/year | L | 2000 | 1.7 | 1.7 | 1.7 | 1.00 | 1.00 | 1.00 | (Walkden-Brown <i>et al.</i> 2012) |
| <i>Farmlet productivity</i> | | | | | | | | | | |
| Liveweight/ha (weaners) | kg/ha | H | 2000 drop | 75.0 | 69.5 | 61.8 | 1.00 | 0.93 | 0.82 | (Hinch <i>et al.</i> 2012a) |

| | | | | | | | | | | |
|----------------------------------|-----------------------|---|-----------|-----------|-----------|---------|-------------|-------------|-------------|-----------------------------|
| Greasy fleece weight/ha | kg/ha/yr | H | 2001 | 12.6 | 11.3 | 10.8 | 1.00 | 0.90 | 0.86 | (Cottle <i>et al.</i> 2012) |
| Stocking rate (annual average) | dse/ha | H | 2000-2001 | 7.5 | 6.7 | 8.2 | 0.92 | 0.82 | 1.00 | (Hinch <i>et al.</i> 2012a) |
| <i>Financial</i> | | | | | | | | | | |
| Fodder cost/farm | \$/year | L | 2000 | \$4,135 | \$2,894 | \$3,791 | 0.70 | 1.00 | 0.76 | (Scott <i>et al.</i> 2012b) |
| Wool value/ewe | \$/hd | H | 2000-2001 | \$31.59 | \$32.66 | \$32.75 | 0.96 | 1.00 | 1.00 | (Cottle <i>et al.</i> 2012) |
| Wool income/ha | \$/hd | H | 2000-2001 | \$182 | \$178 | \$177 | 1.00 | 0.98 | 0.97 | (Cottle <i>et al.</i> 2012) |
| Gross margin/ha (full-size farm) | \$/ha | H | 2000-2001 | \$215 | \$230 | \$181 | 0.93 | 1.00 | 0.79 | (Scott <i>et al.</i> 2012b) |
| Cash position (full-size farm) | \$/farm | H | 2000-2001 | -\$76,393 | \$118,908 | \$2,486 | 0.00 | 1.00 | 0.40 | (Scott <i>et al.</i> 2012b) |
| Equity | % | H | 2000-2001 | 92% | 100% | 100% | 0.92 | 1.00 | 1.00 | (Scott <i>et al.</i> 2012b) |
| Labour per dse | hours/year/ dse/ha | L | 2000-2001 | 35 | 32 | 43 | 0.91 | 1.00 | 0.74 | (Scott <i>et al.</i> 2012c) |
| Average index | | | | | | | 0.91 | 0.86 | 0.86 | |

Table 2. Average data from farmlets A, B and C late in the trial (2003 to 2006) for a wide array of soil, pasture, ground cover, livestock production, animal health, farmlet productivity and financial measurements and proportions of the highest data value of each measurement, normalised on a 0-1 scale.

| Factor | Units | Aim (High/Low) | Year(s) | Average data | | | Proportion of maximum value (for aim of H)/ minimum (for aim of L) | | | Source of data from related publication |
|----------------------------------|--------------|-------------------|-----------|--------------|------|------|---|------|------|--|
| | | | | A | B | C | A | B | C | |
| <i>Soil layer</i> | | | | | | | | | | |
| Soil-N | mg N/kg soil | H | 2003-2006 | 24.9 | 14.1 | 10.4 | 1.00 | 0.57 | 0.42 | (Guppy <i>et al.</i> 2012) |
| Soil-P | mg P/kg soil | H | 2003-2006 | 58.1 | 24.1 | 30.1 | 1.00 | 0.41 | 0.52 | (Guppy <i>et al.</i> 2012) |
| Soil-S | mg S/kg soil | H | 2003-2006 | 12.2 | 6.4 | 6.8 | 1.00 | 0.52 | 0.56 | (Guppy <i>et al.</i> 2012) |
| <i>Pasture layer</i> | | | | | | | | | | |
| Sown perennial grasses | % | H | 2005-2006 | 47.3 | 17.4 | 26.7 | 1.00 | 0.37 | 0.57 | (Shakhane <i>et al.</i> 2012b) |
| Legume | % | H | 2005-2006 | 5.3 | 1.3 | 1.6 | 1.00 | 0.24 | 0.30 | (Shakhane <i>et al.</i> 2012b) |
| Warm season grasses | % | L | 2005-2006 | 15.7 | 29.6 | 29.9 | 1.00 | 0.53 | 0.52 | (Shakhane <i>et al.</i> 2012b) |
| Green dry matter | kg DM/ha | H | 2005-2006 | 942 | 905 | 989 | 0.95 | 0.91 | 1.00 | (Shakhane <i>et al.</i> 2012a) |
| Digestibility of green | % | H | 2005-2006 | 66 | 55 | 58 | 1.00 | 0.84 | 0.87 | (Shakhane <i>et al.</i> 2012a) |
| Dead dry matter | kg DM/ha | L | 2005-2006 | 1048 | 2131 | 2055 | 1.00 | 0.49 | 0.51 | (Shakhane <i>et al.</i> 2012a) |
| Legume dry matter | kg DM/ha | H | 2005-2006 | 51 | 5 | 6 | 1.00 | 0.10 | 0.12 | (Shakhane <i>et al.</i> 2012a) |
| Greenness (satellite NDVI) | NDVI | H | | 554 | 481 | 496 | 1.00 | 0.87 | 0.90 | (Donald <i>et al.</i> 2012) |
| <i>Sustainability</i> | | | | | | | | | | |
| Ground cover | % | H | 2005-2006 | 88 | 96 | 98 | 0.90 | 0.98 | 1.00 | (Shakhane <i>et al.</i> 2012a) |
| <i>Livestock production</i> | | | | | | | | | | |
| Greasy fleece weight/head (ewes) | kg/hd | H | 2003-2005 | 3.6 | 3.5 | 3.1 | 1.00 | 0.97 | 0.86 | (Cottle <i>et al.</i> 2012) |
| Fibre diameter (ewes) | microns | L | 2003-2005 | 18.7 | 18.5 | 18.2 | 0.97 | 0.98 | 1.00 | (Cottle <i>et al.</i> 2012) |
| Staple length (ewes) | mm | H | 2003-2005 | 88.8 | 86.1 | 80.9 | 1.00 | 0.97 | 0.91 | (Cottle <i>et al.</i> 2012) |
| Staple strength (ewes) | N/ktex | H | 2003-2005 | 42.0 | 40.2 | 43.9 | 0.96 | 0.92 | 1.00 | (Cottle <i>et al.</i> 2012) |
| Liveweight/head (ewes) | kg/hd | H | 2004-2005 | 46.3 | 44.6 | 42.6 | 1.00 | 0.96 | 0.92 | (Hinch <i>et al.</i> 2012a) |

| | | | | | | | | | | |
|----------------------------------|-----------------------|-----------------------|-----------|------------|-----------|-----------|-------------|-------------|-------------|------------------------------------|
| Fat score/head (ewes) | score (1-5) | H | 2004-2005 | 2.7 | 2.5 | 2.5 | 1.00 | 0.90 | 0.91 | (Hinch <i>et al.</i> 2012b) |
| Pregnancy scan | proportion | H | 2005 | 0.98 | 0.81 | 0.82 | 1.00 | 0.83 | 0.84 | (Hinch <i>et al.</i> 2012b) |
| Lambs marked/ewe joined | % | H | 2004 | 85 | 90 | 84 | 0.94 | 1.00 | 0.93 | (Hinch <i>et al.</i> 2012b) |
| Lamb mortality (scan to weaning) | % | L | | 31 | 17 | 19 | 0.55 | 1.00 | 0.89 | (Hinch <i>et al.</i> 2012b) |
| Liveweight/head (weaners) | kg/hd | H | 2004 drop | 18.0 | 20.0 | 19.6 | 0.90 | 1.00 | 0.98 | (Hinch <i>et al.</i> 2012a) |
| Liveweight gain/head (cattle) | kg/hd/day | H | 2004-2005 | 0.97 | 0.95 | 0.89 | 1.00 | 0.98 | 0.92 | (Hinch <i>et al.</i> 2012a) |
| Parasites (faecal egg count) | number/gram | L | 2002-2006 | 594.4 | 790.2 | 367.9 | 0.62 | 0.47 | 1.00 | (Walkden-Brown <i>et al.</i> 2012) |
| Number of drenches/mob/year | number/year | L | 2005-2006 | 2.7 | 3.0 | 1.9 | 0.70 | 0.62 | 1.00 | (Walkden-Brown <i>et al.</i> 2012) |
| <i>Farmlet productivity</i> | | | | | | | | | | |
| Liveweight/ha (weaners) | kg/ha | H | 2004 drop | 89.7 | 64.0 | 54.7 | 1.00 | 0.71 | 0.61 | (Hinch <i>et al.</i> 2012a) |
| Greasy fleece weight/ha | kg/ha/yr | H | 2004-2005 | 23.4 | 17.1 | 13.5 | 1.00 | 0.73 | 0.57 | (Cottle <i>et al.</i> 2012) |
| Liveweight gain/ha (cattle) | kg/ha/yr | H | 2004-2005 | 84.49 | 71.40 | 78.72 | 1.00 | 0.85 | 0.93 | (Hinch <i>et al.</i> 2012a) |
| Stocking rate (annual average) | dse/ha | H | 2004-2005 | 11.9 | 8.2 | 7.6 | 1.00 | 0.69 | 0.63 | (Hinch <i>et al.</i> 2012a) |
| Silage produced | t Digestible DM/ha | H | 2005 | 20.2 | 0.0 | 0.0 | 1.00 | 0.00 | 0.00 | (Shakhane <i>et al.</i> 2012a) |
| <i>Financial</i> | | | | | | | | | | |
| Fodder cost/farmlet | \$/year | L | 2005 | \$13,761 | \$8,247 | \$7,448 | 0.54 | 0.90 | 1.00 | (Scott <i>et al.</i> 2012b) |
| Wool value/ewe | \$/hd | H | 2003-2005 | \$39.80 | \$41.05 | \$38.06 | 0.97 | 1.00 | 0.93 | (Cottle <i>et al.</i> 2012) |
| Wool income/ha | \$/ha | H | 2003-2005 | \$303 | \$215 | \$180 | 1.00 | 0.71 | 0.59 | (Cottle <i>et al.</i> 2012) |
| Gross margin/ha (full-size farm) | \$/ha | H | 2003-2005 | \$303 | \$241 | \$214 | 1.00 | 0.79 | 0.71 | (Scott <i>et al.</i> 2012b) |
| Cash position (full-size farm) | \$/farm | H | 2003-2005 | -\$152,316 | \$461,125 | \$241,595 | 0.00 | 1.00 | 0.64 | (Scott <i>et al.</i> 2012b) |
| Equity | % | H | 2003-2005 | 92% | 100% | 100% | 0.92 | 1.00 | 1.00 | (Scott <i>et al.</i> 2012b) |
| Labour per dse | hours/year/ dse/ha | hours/year/ dse/ha | 2005 | 18 | 20 | 22 | 1.00 | 0.90 | 0.82 | (Scott <i>et al.</i> 2012c) |
| Average index | | | | | | | 0.91 | 0.76 | 0.76 | |




Figure 1. Diagram showing relativity between farmlets A, B and C early in the trial (from 2000 to 2001) for a wide array of soil, pasture, ground cover, livestock production, animal health, farmlet productivity and financial measurements, normalised to a 0-1 scale (proportions extracted from Table 1).




Figure 2. Diagram showing relativity between farmlets A, B and C late in the trial (from 2003 to 2006) for a wide array of soil, pasture, ground cover, livestock production, animal health, farmlet productivity and financial measurements, normalised to a 0-1 scale (proportions extracted from Table 2).