



Growing Australia's Future

Username  Password    
[Sign up now!](#) | [Forgotten your password?](#) [help](#)

## Australian Agronomy Conference

[1980 1st AAC](#)
[1982 2nd AAC](#)
[1985 3rd AAC](#)
[1987 4th AAC](#)
[1989 5th AAC](#)
[ASA 1992](#)
[1993 7th AAC](#)
[1996 8th AAC](#)
[1998 9th AAC](#)
[2001 10th AAC](#)
[2003 11th AAC](#)
[2004 12th AAC, 4th ICSC](#)
[2006 13th AAC](#)
[2008 14th AAC](#)
[2010 15th AAC](#)
[2012 16th AAC](#)
[Agriculture](#)
[2012 Authors](#)
[Breeding](#)
[Climate change](#)
[Crop Development](#)
[Crop production](#)
[Disease](#)
[Nutrition](#)
[Pastures](#)
[Seasonal production of coloured brome \(\*Bromus coloratus\* Steud\) cv. Exceltas, a new high quality perennial temperate pasture grass](#)
[Austroanthonia spp. in grazed native pastures on the Monaro tableland, NSW](#)
[Southern Australian feed-base pasture audit](#)
[Use of non fertiliser growth promotants as alternatives to urea on high production pastures](#)
[Pasture legume production severely reduced when co-sown with winter forage cereals](#)
[Hard seed breakdown rates of six tropical ley](#)
[Home](#) » [Publications](#) » [Australian Agronomy Conference](#) » [2012 16th AAC](#) » [Pastures](#)
 Print
 [← Previous](#)
 Menu
 [Next →](#)

# To under-sow or not? A decision support tool to determine the most profitable method of pasture establishment

Jeff McCormick<sup>1,3</sup>, Richard Hayes<sup>1,3</sup>, Guangdi Li<sup>1,3</sup>, Tom Nordblom<sup>1,3</sup>, Geoff Casburn<sup>1,3</sup>, Tim Hutchings<sup>1</sup>, Andrew D. Moore<sup>2,3</sup>, Eric Zurcher<sup>2,3</sup>, Mark Peoples<sup>2,3</sup> and Tony Swan<sup>2,3</sup>

<sup>1</sup> EH Graham Centre for Agricultural Innovation (an alliance between NSW Department of Primary Industries and Charles Sturt University), Wagga Wagga Agricultural Institute, Pine Gully Rd, Wagga Wagga NSW 2650, [richard.hayes@dpi.nsw.gov.au](mailto:richard.hayes@dpi.nsw.gov.au)

<sup>2</sup>CSIRO Sustainable Agriculture Flagship, GPO Box 1600, Canberra ACT 2601

<sup>3</sup>CRC for Future Farm Industries, 35 Stirling Highway, Crawley, WA 6009, Australia

## Abstract

A new decision support tool (DST) is under development to enable users to determine whether it is more profitable to establish a perennial pasture alone or under-sowing with cover-crops in the uniform rainfall zone of southern NSW (450-650 mm). The DST operates under the premise that a pasture is to be sown in a particular paddock in the coming year. A simple underlying model focuses on the profitability of each pasture sowing method at the end of the pasture phase, including all variable costs and income. The DST enables the user to input a range of information including costs of establishment and income from both grain and livestock. Exploratory analyses with the under-sowing tool for the uniform rainfall zone of southern NSW show that pastures established by under-sowing can be more profitable under different scenarios even when only half as productive as a straight-sown pasture due to the returns from the cover-crop. Pastures established by under-sowing tend to be more profitable when the grain yield is greater than 2.5 T/ha or grain price is greater than \$180/T. Pastures sown without a cover crop were more profitable when there was the potential for high stocking rates with high stock returns and a longer pasture phase (> 4 years). Future developments of the model will seek to include climate variability for specific sites to enable producers to determine the risk of under-sowing in their locality.

## Key Words

Cover-crops, gross margin, mixed farming system

## Introduction

Pasture establishment in the cropping zone has commonly occurred by sowing the pasture species under a cover crop. Research and extension information on under-sowing is available from the last 80 years with many recommending that under-sowing should not be used to establish pastures due to a higher rate of failure and less productive pastures (Moodie 1936; Smith and Argyle 1964; Peart and Scott 1969; Cregan 1985; Dear 1986). Considering this, a recent survey has revealed that within the uniform rainfall cropping zone of southern NSW that 83% of farmers regularly under sow (Li *et al.* 2010). Generally farmers use under-sowing because the grain yield covers the cost of sowing the pasture. Most research has focused on pasture density and biomass production of the different establishment methods but has not sought to demonstrate increased livestock productivity that could outweigh income from grain production due to a cover crop. This DST seeks to bring the costs and incomes for the pasture phase to inform producers whether under-sowing is the right method of establishment for them.

## Methods

### Model development

The DST has been created in an MS Excel file with a user-friendly display where users can input their own data and determine which sensitivity graphs

Search Papers

 all
  journal
  2012



## Publications

Browse our full text  
[\[Conference Publications.....\]](#)



pasture legumes.

Potential of *Desmanthus* spp. in northern New South Wales

Companion legume species maximise productivity of chicory (*Cichorium intybus*)

Success of perennial pasture establishment at different sowing times and under a cover crop in the mixed farming zone

Summer sowing: a new alternative technique to introduce annual pasture legumes into mixed farming systems

Seed rain of *Microlaena stipoides*

Novel Cocksfoots for SE Australia &#8211; Establishment and production

A survey of land use and management, North-West Slopes of New South Wales

The germination, passage and viability of *Desmanthus virgatus* (L.) Willenow seed through sheep and its implication for dispersal in tropical rangelands

Low-input, high-quality legume hays for north Queensland.

Evaluating new *phalaris* populations for lower rainfall margins in SE Australia

The advantages and disadvantages of pasture cropping in Western Australia

The search for sustainable stocking rates on Chinese desert steppe

Improving mineral availability for grazing livestock in Australian pasture systems by using plantain and lucerne

A survey of farmer practice on the establishment, duration and production of pastures in the mixed farming zone of southern New South Wales

To under-sow or not? A decision support tool to determine the most profitable method of

they want to be displayed.

The underlying calculation for the DST is the net income from under-sowing (US) pasture establishment method minus the net income from straight-sowing (SS) the pasture for the length of the pasture phase (eqn 1).

$$(\text{Grain income} + \text{US stock income} - \text{US variable cost}) - (\text{SS stock income} - \text{SS variable cost}) \text{ eqn 1.}$$

Where

$$\text{Grain income} = \text{grain yield} \times \text{grain price}$$

$$\text{US stock income} = \text{stocking rate} \times \$/\text{DSE} \times (\text{pasture years} - 1) \times \text{US relative effect}$$

$$\text{SS stock income} = \text{stocking rate} \times \$/\text{DSE} \times (\text{pasture years} - 1)$$

$$\text{US variable cost} = \text{cost of establishing grain crop and pasture by under-sowing}$$

$$\text{SS variable cost} = \text{cost of establishing straight sown pasture}$$

The value for \$/DSE has been determined from NSW DPI budgets and is the net income from livestock and includes costs for stock and pasture management. The years are for the length of the intended pasture phase minus the establishment year where grazing is limited. The DST does not calculate pasture production *per se* but rather calculates the differences in stocking rates for the different establishment options, which we assume is related to pasture production. The under-sowing relative effect is the proportion of production of an under-sown pasture compared to a straight sown pasture with 0 being no production and 1 being equal production.

## Model inputs

An important component of the model is the capacity for the user to change a range of inputs to match their enterprise. The inputs in the DST include expected grain yield, grain price, stocking rate and stock earnings, establishment costs, the length of the pasture phase and relative effect that under-sowing has on pasture production.

**Table 1. Parameters and input data for Decision Support Tool**

<b>Crop Income</b>	Grain Price	\$180/T
	Grain Yield	2.5 T/ha
<b>Livestock Income</b>	Stocking Rate	10 DSE/Ha
	\$/DSE	\$25/DSE
<b>Costs</b>	US establishment cost	\$200/Ha
	SS establishment cost	\$120/Ha
<b>Years of Pasture</b>	Years	4
<b>Reduced pasture production due to under-sow</b>	US relative effect (0-1)	0.5

## Model outputs

The DST provides a single number to estimate which method of pasture establishment is more profitable. If the returned value is positive then greater profitability is obtained from under-sowing. In comparison if the value is negative, directly sowing the pasture would be more profitable. A number of sensitivity graphs are provided to determine what factors change the result.

## Results

Using the values in Table 1 applied in equation 1 gives a value of -\$5.00 which indicates that straight sowing the pasture is more profitable for the length of the pasture phase (Figure 1). The DST comments that this value is "too close to call" and for the user to determine which will be more profitable. With the costs in Table 1, the point of equilibrium for the sowing methods in relation to grain yield is at 2.5 t/ha (Figure 2a). Higher grain yield would lead to under-sowing being more profitable than straight sowing the pasture. Similarly, an increase in grain price will lead to under-sowing

pasture establishment

Nutritive value of silage from perennial ryegrass, plantain and lucerne pastures in south west Victoria

Crop yield, pasture yield, and environmental impact of pasture cropping with sub-tropical perennials.

Identifying lucerne growers' usage and needs through a survey for breeding new generation lucerne varieties

Opportunities for grazing crop/pasture intercrops

Weed control options for pasture cropping systems in the West Midlands of Western Australia

Evaluating the importance of a potential source of error when applying shoot 15N labelling techniques to legumes to quantify the below-ground transfer of nitrogen to other species

Important principles of pasture management demonstrated within the Cicerone Project's grazed farmlet experiment: some personal reflections

Tasmanian pasture resource audit: Snapshot of functional group composition in 2011

Increasing soil fertility and altering legume-grass balance as adaptation strategies for sustainable livestock production under climate change

Graziers, pasture seed industry and researchers are concerned about pasture productivity decline.

New tall fescue cultivars for medium to low rainfall environments in southern Australia

Assessment of Lotus


becoming the most profitable option (Figure 2b). Increases in stocking rate or stock price will favour establishment of pastures by straight sowing (Figure 2c, d). The length of the pasture phase is a strong determinant of profitability for each sowing method with pasture phases greater than four years tending to be more profitable with straight sowing (Figure 2e). The under-sowing relative factor was highly significant in determining the success of each sowing method (Figure 2f). Increasing this relative factor leads to increases in profitability of the under-sowing method.

## Discussion

The primary purpose of the DST is to enable users to include all necessary inputs into the decision making process on whether to establish pasture by the under-sowing method. This model allows users to input their own data to determine what is more profitable. The graph on the user interface enables the user to determine how sensitive each parameter is to variation.

There are some difficulties with the DST. It is not a biological model as it does not predict pasture production for a certain set of parameters. This is particularly important for determining the value of the US relative value. It is likely there would be different perceptions of the under-sowing effect but the limited data tends to indicate that the reduction in pasture growth results in a relative value of between 0.5-0.8. This can have a large influence on the DST. The other difficulty regarding the model is in relation to pasture establishment failure. The model does not determine whether there has been a pasture failure. Currently there is no published data that recommends when pasture failure has occurred or when pastures should be re-sown.

The DST is currently being showcased to producers to determine their interest in the model and whether it corresponds to what they observe. It will be important to simplify some inputs such as \$/DSE as these are difficult for the individual farmer to quantify. Incorporating climate data into the model will enable specific sites to determine over a large number of seasonal years whether under-sowing is more profitable or not.



**Figure 1. The user interface of the decision support tool using the input data from the materials and methods (Table 1).**

## Conclusion


The DST enables users to identify whether under-sowing is more profitable for their farming situation. The range of sensitivity graphs helps determine how sensitive a particular parameter is. Grain yields greater than 2.5 t/ha with grain prices greater than \$180/t will tend to improve the profitability of under-sowing. Alternatively increases in stocking rates and prices will result in straight sowing pasture being more profitable. The length of the pasture phase and the cover crop relative effect are also significant factors in the profitability of each pasture establishment method.

tenuis for desirable seed production characteristics

Leucaena: Highlighting the value of good agronomy for establishing pasture systems.

EverFarm: Piloting a strategic decision making process to improve grazing farm systems.

Pests	<input type="checkbox"/>
Plenary papers	<input type="checkbox"/>
Precision Agriculture	<input type="checkbox"/>
Soil Water Management	<input type="checkbox"/>
Weeds	<input type="checkbox"/>



**Figure 2. Sensitivity graphs of total gross margins for the pasture phase for a range of parameters based on original analysis. Parameters include a) cover crop grain yield, b) grain price, c) stocking rate, d) stock price, e) length of pasture phase and f) under-sow relative effect. Solid line – under-sow establishment, dashed line – straight sowing and dotted line – point of intersection.**

## References

Cregan, PD (1985) 'Managing the seedling for establishment, Pasture renovation Proceedings Grasslands Society of Victoria 26th Annual Conference.' Melbourne.

Dear, BS GM Murray, HM Dengate (Eds) (1986) 'Pasture establishment using cover crops - is it worthwhile?', 29th Southern Cereal Improvement Conference.' Wagga Wagga, NSW. (Department of Agriculture:

Li, GD, Dear, BS, Casburn, G, Walker, J, Bowden, P, Hayes, RC (2010) 'A survey of the current and future use of lucerne, chicory and perennial grasses in pasture-crop rotations in southern New South Wales, 15th Australian Agronomy Conference.' Lincoln, New Zealand.

Moodie, AWS (1936) Sowing lucerne with wheat. *The Agricultural Gazette of New South Wales* **47**, 419.

Peart, GR, Scott, RC (1969) Sowing methods for establishment of dryland lucerne. *The Agricultural Gazette of New South Wales* **80**, 66.

Smith, JAC, Argyle, DB (1964) Establishing wheatbelt pastures - don't use a cover-crop. *Journal of Agriculture Western Australia* **5**, 134.

[◀ Previous](#) [▲ Top](#) [Next ▶](#)